

# 8th EOS Topical Meeting on Diffractive Optics 2012 (DO 2012)

27 February - 1 March 2012, Conference Centre of Delft University, Delft, Netherlands

## FINAL PROGRAMME

### Sponsors



### Cooperating organisation



### Media Partner



# EOS

European Optical Society

Annual Meeting

# EOSAM 2012 moves to Aberdeen



Aberdeen Exhibition and Conference Centre, Scotland (UK) | 25 - 28 September 2012

## Present your research in the energy capital of Europe!

### Topical Meetings

- TOM 1 Biophotonics
- TOM 2 Silicon Photonics
- TOM 3 Nanophotonics & Metamaterials
- TOM 4 Micro-Optics
- TOM 5 Organic Photonics & Electronics
- TOM 6 Nonlinear Photonics
- TOM 7 Optical Systems for Energy & Production Industries

### Workshop

Continuing education: Short courses for industry

### Exhibition with special focus on:

- Photonics for offshore applications: blue photonics®
- Biomedical photonics
- Organic optoelectronics
- Micro-optical components and systems

Sign up for updates at [www.myeos.org/events/eosam2012](http://www.myeos.org/events/eosam2012)

**General information**

About EOS .....	1
Venue & Getting there .....	2-3
Accommodation & Alternative hotels in Delft .....	4-5
Information for authors and attendees .....	6-8

**EOS Topical Meeting on Diffractive Optics 2012**

Welcome to DO 2012 .....	9
Programme committee.....	9
Plenary & Invited talks.....	10-11

**Programme**

Daily overview .....	12
----------------------	----

**Oral presentations**

• Monday, 27 February .....	13-16
• Tuesday, 28 February .....	17-19
• Wednesday, 29 February .....	20-22
• Thursday, 1 March .....	23-24

**Poster presentations**

• Poster session .....	25-33
------------------------	-------

Registration fax form .....	34-35
-----------------------------	-------

Upcoming EOS events.....	36
--------------------------	----

## ABOUT EOS

### History

The European Optical Society (EOS) was founded in 1991. The purpose of the society is to contribute to progress in optics and related sciences, and to promote their applications at the European and international levels, by bringing together individuals and legal entities involved in these disciplines and their applications. EOS is a not for profit organisation and serves as the joint forum for all individuals, companies, organisations, educational institutions, and learned and professional societies, who recognise the opportunity and challenge that a common European base provides for the development of optics in its broadest sense. EOS organises recognized topical meetings, conferences, workshops and other events, publishes journals and is an important player on the European level. 22 national optical societies and a great number of individuals and companies are currently members of EOS ([www.myeos.org](http://www.myeos.org)).



### Membership modes and fees

#### Individual membership

Annual fee: 50 €

#### Individual membership through an EOS Branch

Every member of an EOS Branch is automatically an individual member of the EOS, too, with all benefits.

Annual fee: included in the Branch membership fee

#### Student membership

Annual fee: 10 €

#### Associate membership through an EOS Affiliated Society

Every member of an EOS Affiliated Society is automatically an associate member of the EOS, too, but with limited benefits.

Annual fee: included in the Affiliated Society membership fee

### EOS membership - Join us and...

- Be a part of the umbrella organisation of the national optical societies in Europe
- Connect with colleagues from all over Europe and beyond
- Contribute to strengthening Europe's future in optics and photonics
- Stay up-to-date about European Research Funding
- Benefit from discounts on EOS events and publications in the EOS online journal JEOS:RP
- Receive the Annual EOS Member Directory - your guide to the European optics and photonics community

### Activities

- Organisation of topical meetings, workshops and conferences, and endorsement of other scientific events
- Operation of a virtual platform for the European optics and photonics community at [www.myeos.org](http://www.myeos.org)
- Focus Groups and Student Clubs
- Publication of JEOS:RP, the electronic Journal of the European Optical Society - Rapid Publications ([www.jeos.org](http://www.jeos.org))
- EOS Print Newsletter to be released three-monthly in cooperation with AT-Fachverlag
- Representation of the optics and photonics community on the European level (Photonics21 Technology Platform)
- Annual award of the EOS Prize

### Upgrade for associate members

Upgrade to an individual EOS membership with full benefits.  
Annual fee: 12.50 €

### Corporate membership through an EOS Branch or Affiliated Society

Annual fee: 200 €

### Direct corporate membership

Annual fee: 300 €

### How to join?

To join the EOS as an individual, student or corporate member, please see our website at [www.myeos.org/members](http://www.myeos.org/members).

### Questions?

Please contact the EOS office at [info@myeos.org](mailto:info@myeos.org).


## VENUE

Discover one of the oldest cities in the Netherlands called *de Prinsenvstad*, the Princes City, since William of Orange held court in Delft in the 16th century. Stroll through the well-preserved, lively historical centre, with characteristic canals, ancient merchant houses, old churches and the splendid city hall which make Delft a perfect location for a high-quality EOS Topical Meeting in an enjoyable atmosphere. See also: [www.delft.nl](http://www.delft.nl)



### The 8th EOS Topical Meeting on Diffractive Optics (DO2012)

takes place at the Conference Centre of the Delft University of Technology, located in the heart of the University campus and in walking distance from the Delft city centre and public transport facilities.



Conference Centre of the  
Delft University of Technology  
Mekelweg 5  
2628 CC Delft, Netherlands

phone +31 (0)15 - 27 88022  
fax +31 (0)15 - 27 86755

## GETTING THERE

### BY PLANE

The *Amsterdam Airport Schiphol* that can be reached from various international airports in the world is about 45 min away by train. For Delft, you have to change at *Leiden Centraal* or *Den Hague Hollands Spoor (Den Haag HS)*. If you prearrange a taxi to pick you up at Schiphol, expect to pay about € 60 to Delft.

The *Rotterdam The Hague Airport* is only 12 kilometres away and can be reached by RET city bus 33 that runs frequently to *Rotterdam Centraal Station* in about 20 min. From there you can go to Delft by train in 20 minutes. When taking an airport taxi from *Rotterdam The Hague Airport*, expect to pay about € 35 for the trip to Delft.

### BY TRAIN

Delft has two railway stations: *Delft* (near the city center and best for the university) and *Delft Zuid* (Delft's southern residential area). Trains run from:

*Den Haag Central Station* or *Hollands Spoor* (10 min, € 2.20)

*Amsterdam* (1 h, € 11.50)

*Rotterdam* (15 min, € 2.90)

*Schiphol Airport* (45 min, € 8.70)

(Please note that the train from Schiphol does not stop at Delft and you need to change at Leiden or *Den Haag HS*)

For the journey planner please refer to [www.ns.nl/en](http://www.ns.nl/en)

### BY TRAM

From *Den Haag HS* also tramline 1 runs to Delft. Travelling time is approximately 20 minutes and a one-way ticket costs € 1.74.

On the trams you must use the OV-Chipcard or a day-pass which can be bought at the ticket office and vending machines at the station.

More information at [www.htm.net](http://www.htm.net) and at the journey planner at <http://journeyplanner.9292.nl/Default.aspx>

### BY CAR

Coming from Amsterdam or Schiphol, Delft is at the end of motorway A4. Coming from Rotterdam, Delft has three exits off the A13 motorway. The city centre is best reached via the IKEA exit or exit Noord.

## GETTING THERE - continued

### BY CAR

#### GETTING AROUND IN DELFT

During the next years construction works are in progress as a new train tunnel is being built through Delft. Therefore the flow of traffic and parking is severely affected and it is recommended not to use the car while visiting Delft.

Anyone who still goes by car please note that in the TU Delft area parking is possible for free but not in the old city. It has been made largely car-free but there are three large car parks at the edge of the center (*Zuidpoort garage*, *Koepoort garage* and *Phoenix garage*). These parking garages are opened 24 hours a day and parking fee is € 2.00 per hour.

Download more information about parking in Delft and a map with all possible spots at [www.myeos.org/download/DO2012/Folder%20Parkeren%20in%20het%20centrum%20van%20Delft.pdf](http://www.myeos.org/download/DO2012/Folder%20Parkeren%20in%20het%20centrum%20van%20Delft.pdf).

Note also that parking policy differs per area and is not always clear. Beware of signs "alleen vergunninghouders" (permit holders only) in some streets especially in residential areas. Sometimes only a floor tile with a "P" is visible. This means permit parking only. Unauthorized parking will cost about € 70.

### BY BUS

From the *Station Centrum*, which is at the back of Delft railway station, several bus lines run to the station *Aula TU*.

51 (direction: *Zoetermeer, Centrum West*)

Runs only Monday-Friday. Timetable at [www.veolia-](http://www.veolia-transport.nl/netherlands-transport/ressources/documents/2/12074,201010_haagl_02_www_redu.pdf)

[transport.nl/netherlands-transport/ressources/documents/2/12074,201010\\_haagl\\_02\\_www\\_redu.pdf](http://transport.nl/netherlands-transport/ressources/documents/2/12074,201010_haagl_02_www_redu.pdf).

69 (direction: *Delft, Heertjeslaan*)

Runs only Monday-Friday. Timetable at [www.veolia-transport.nl/netherlands-transport/ressources/documents/2/12077,201010\\_haagl\\_04\\_www\\_redu.pdf](http://www.veolia-transport.nl/netherlands-transport/ressources/documents/2/12077,201010_haagl_04_www_redu.pdf).

121 (direction: *Zoetermeer, Centrum West*)

Timetable at [www.veolia-transport.nl/netherlands-transport/ressources/documents/2/12074,201010\\_haagl\\_02\\_www\\_redu.pdf](http://www.veolia-transport.nl/netherlands-transport/ressources/documents/2/12074,201010_haagl_02_www_redu.pdf).

174 (direction: *Zoetermeer, Centrum West*)

Timetable at [www.qbuzz.nl/regio+rotterdam/route+tijden/dienstregeling/cDU410\\_Dienstregeling.aspx?intBLDId=1904&intDKId=3486&intStopId=15744&strHours=Schedule\\_2](http://www.qbuzz.nl/regio+rotterdam/route+tijden/dienstregeling/cDU410_Dienstregeling.aspx?intBLDId=1904&intDKId=3486&intStopId=15744&strHours=Schedule_2).

See the network map of Delft at [www.veolia-transport.nl/netherlands-transport/ressources/documents/2/15128,Delft\\_Abrikaart\\_2010\\_2011\\_v3.pdf](http://www.veolia-transport.nl/netherlands-transport/ressources/documents/2/15128,Delft_Abrikaart_2010_2011_v3.pdf).

### BY FOOT

The city centre of Delft is small enough to walk across in 10 minutes. Most streets in the city centre are pedestrianised and only taxis are allowed to use these streets, too.

The walking time from the city center to the aula of TU Delft is approximately 10 -15 minutes.

### FURTHER INFORMATION

More information about Delft at [www.delft.nl/delften](http://www.delft.nl/delften).



## ACCOMODATION

Rooms at special rates have been blocked at four selected hotels. Find the booking details below.

**HAMPSHIRE Hotel Delft Centre \*\*\*\***

Phone: +31 (0)15 212 21 25

Fax: +31 (0)15 212 21 25

Koepoortplaats 3  
2612RR Delft, NL

[reservations@hoteldelftcentre.nl](mailto:reservations@hoteldelftcentre.nl)

[www.hoteldelftcentre.nl](http://www.hoteldelftcentre.nl)

Remarks: Free Internet available in the rooms, free Wi-Fi in common areas. Discount tickets for the parking garage Koepoort available, € 10.00 per 24 hours.

Rate(s): Room for single occupancy incl. breakfast € 99.50 plus visitors' tax.

Keyword: Please use this [registration form](#) when making your reservation and mention the keyword "EOS Conference".

This offer was valid until 31 January 2012. Afterwards a reservation at this rate is possible only upon availability.

**Hotel Johannes Vermeer \*\*\***

Phone: +31 (0)15 212 64 66

Fax: +31 (0)15 213 48 35

Molslaan 20  
2611 RM Delft, NL

[info@hotelvermeer.nl](mailto:info@hotelvermeer.nl)

[www.hotelvermeer.nl](http://www.hotelvermeer.nl)

Remarks: Free Wi-Fi in the rooms included. Discount tickets for the parking garage Zuidpoort available, € 10.00 per 24 hours.

Rate(s): Room for single occupancy incl. breakfast € 106.40 plus visitors' tax.

Room for double occupancy incl. breakfast € 118.75 plus visitors' tax.

Keyword: Please mention "EOS Conference" when making your reservation.

This offer was valid until 15 January 2012.

**Hotel Juliana \*\*\***

Phone: +31 (0)15 256 76 12

Fax: +31 (0)15 256 57 07

Trompstraat 33  
2628 RC Delft, NL

[info@hoteljuliana.nl](mailto:info@hoteljuliana.nl)

[www.hoteljuliana.nl](http://www.hoteljuliana.nl)

Remarks: Free Wi-Fi in the rooms and free parking included.

Rate(s): Single room incl. breakfast € 79.80 plus visitors' tax.

Double room for single occupancy incl. breakfast € 89.30 plus visitors' tax.

Double room for double occupancy incl. breakfast € 98.80 plus visitors' tax.

Keyword: Please mention "EOS Conference" when making your reservation.

This offer was valid until 31 January 2012. Afterwards a reservation at standard rates is possible only upon availability.

**Hotel de Plataan \*\*\***

Phone: +31 (0)15 212 60 46

Fax: +31 (0)15 215 73 27

Doelenplein 10  
2611 BP Delft, NL

[info@hoteldeplataan.nl](mailto:info@hoteldeplataan.nl)

[www.hoteldeplataan.nl](http://www.hoteldeplataan.nl)

Remarks: Free Wi-Fi in the rooms and free parking included.

Rate(s): Room for single occupancy incl. breakfast € 89.10 plus visitors' tax.

Room for double occupancy incl. breakfast € 102.60 plus visitors' tax.

Room for three persons incl. breakfast € 143.10 plus visitors' tax.

Room for four persons incl. breakfast € 183,60 plus visitors' tax.

Keyword: Please mention "EOS Conference" when making your reservation.

This offer was valid until 31 January 2012.

## ALTERNATIVE HOTELS IN DELFT

**Please note** that room rates as well as information on internet facilities (internet plugs, Wi-Fi etc.) are taken from the home-pages of the listed hotels. Rates may vary from the prices listed below (e.g. during fairs) according to room availability and reservation date. **Please contact the hotel directly to make your reservation.**

### Hotel de Ark \*\*\*\*

Prices: € 118 - 145 (incl. breakfast) plus visitors' tax  
 Address: Koornmarkt 65, 2611 EC Delft, NL  
 URL: [www.deark.nl](http://www.deark.nl)  
 Contact: [hotel@deark.nl](mailto:hotel@deark.nl)  
 Phone: +31 (0)15 215 79 99  
 Fax: +31 (0)15 214 49 97  
 Remarks: No information about internet access available, parking chargeable

### Hotel Best Western Museumhotels Delft \*\*\*\*

Prices: € 112 - 226 plus visitors' tax  
 (No information about breakfast available)  
 Address: Oude Delft 189, 2611 HD Delft, NL  
 URL: [www.museumhotels.nl](http://www.museumhotels.nl)  
 E-Mail: [info@museumhotels.nl](mailto:info@museumhotels.nl)  
 Phone: +31 (0)15 215 30 70  
 Fax: +31 (0)15 215 30 79  
 Remarks: Free internet access via TV with wireless key board, parking chargeable

### Bridges House Hotel \*\*\*

Prices: € 90 - 200 plus visitors' tax  
 (No information about breakfast available)  
 Address: Oude Delft 74, 2611 CD Delft, NL  
 URL: [www.bridges-house.com](http://www.bridges-house.com)  
 E-Mail: [info@bridges-house.com](mailto:info@bridges-house.com)  
 Phone: +31 (0)15 212 40 36  
 Fax: +31 (0)15 213 36 00  
 Remarks: Free internet access in rooms, parking chargeable

### Campanile Hotel Delft \*\*\*

Prices: € 60 - 100 plus visitors' tax  
 (No information about breakfast available)  
 Address: Kleveringweg 53, 2616 LZ Delft, NL  
 URL: [www.campanile-delft.nl](http://www.campanile-delft.nl)  
 E-Mail: [delft@campanile.com](mailto:delft@campanile.com)  
 Phone: +31 (0)15 279 90 00  
 Fax: +31 (0)15 279 90 01  
 Remarks: Free Wi-Fi access in rooms, free parking

### Hotel Coen Delft \*\*\*

Prices: € 72 - 110 plus breakfast & visitors' tax  
 Address: Coenderstraat 47, 2613 SN Delft, NL  
 URL: [www.hotelcoendelft.nl](http://www.hotelcoendelft.nl)  
 E-Mail: [info@hotelcoendelft.nl](mailto:info@hotelcoendelft.nl)  
 Phone: +31 (0)15 214 59 14  
 Fax: +31 (0)15 212 63 84  
 Remarks: Free Wi-Fi access in rooms, parking chargeable

### Hotel de Emauspoort \*\*\*

Prices: € 85 - 175 (incl. breakfast) plus visitors' tax  
 Address: Vrouwenregt 9, 2611 KK Delft, NL  
 URL: [www.emauspoort.nl](http://www.emauspoort.nl)  
 E-Mail: [info@emauspoort.nl](mailto:info@emauspoort.nl)  
 Phone: +31 (0)15 219 02 19  
 Fax: +31 (0)15 214 82 51  
 Remarks: Free Wi-Fi access in rooms, parking chargeable

### Hotel de Koophandel \*\*\*

Prices: € 94 - 144 plus breakfast & visitors' tax  
 Address: Beestenmarkt 30, 2611 GC Delft, NL  
 URL: [www.hoteldekoophandel.nl](http://www.hoteldekoophandel.nl)  
 E-Mail: [info@hoteldekoophandel.nl](mailto:info@hoteldekoophandel.nl)  
 Phone: +31 (0)15 214 23 02  
 Fax: +31 (0)15 212 06 74  
 Remarks: Chargeable Wi-Fi access in rooms, parking chargeable

### Hotel de Vlaming \*\*\*

Prices: € 70 - 115 plus visitors' tax  
 (No information about breakfast available)  
 Address: Vlamingstraat 52, 2611 KZ Delft, NL  
 URL: [www.hoteldevlaming.nl](http://www.hoteldevlaming.nl)  
 E-Mail: [info@hoteldevlaming.nl](mailto:info@hoteldevlaming.nl)  
 Phone: +31 (0)15 213 21 27  
 Fax: +31 (0)15 212 20 06  
 Remarks: Free internet access in rooms, parking chargeable

### Hotel Leeuwenbrug \*\*\*

Prices: € 83 - 125 (incl. breakfast) plus visitors' tax  
 Address: Koornmarkt 16, 2611 EE Delft, NL  
 URL: [www.leeuwenbrug.nl](http://www.leeuwenbrug.nl)  
 E-Mail: [sales@leeuwenbrug.nl](mailto:sales@leeuwenbrug.nl)  
 Phone: +31 (0)15 214 77 41  
 Fax: +31 (0)15 215 97 59  
 Remarks: No information about internet access available, parking chargeable

### Het Koningshuys \*\*

Prices: € 52 - 99 (incl. visitors' tax) plus breakfast  
 Address: Markt 38-42, 2611 GV Delft, NL  
 URL: [www.hetkoningshuys.eu](http://www.hetkoningshuys.eu)  
 E-Mail: [hotel@hetkoningshuys.eu](mailto:hotel@hetkoningshuys.eu)  
 Phone: +31 (0)15 212 51 15  
 Fax: +31 (0)15 213 60 69  
 Remarks: Internet access available, parking chargeable

### Jorplace First Hostel

Prices: € 19 - 67 (incl. breakfast) plus visitors' tax  
 Address: Voldersgracht 16 - 18, 2611 GV Delft, NL  
 URL: <http://jorplace.nl/delft>  
 E-Mail: [delft@jorplace.nl](mailto:delft@jorplace.nl)  
 Phone: +31 (0)15 570 88 75  
 Remarks: Wi-Fi access available

Further accommodation facilities in Delft are available at: [www.delft.nl/delften/Tourists/Accommodations/Hotels\\_in\\_Delft](http://www.delft.nl/delften/Tourists/Accommodations/Hotels_in_Delft)



## INFORMATION FOR AUTHORS AND ATTENDEES

### EOS REGISTRATION DESK

#### Opening times

Sunday, 26 February	16:00 - 18:00
Monday, 27 February	07:30 - 18:00
Tuesday, 28 February	08:00 - 18:00
Wednesday, 29 February	08:00 - 16:00
Thursday, 1 March	08:00 - 12:00

### INFORMATION / RECEIPTS / CONFIRMATION OF ATTENDANCE / CASH PAYMENT

Payment receipts and confirmation of attendance will be available on-site at the registration desk.

Attendees paying by cash are requested to have the **exact change in Euro**.

### REGISTRATION & FEES

At least one author of an accepted presentation is requested to register properly in advance to the conference.

The full-time-registration fee includes the participation in all four meeting days, one copy of the Topical Meeting digest CD-ROM, coffee breaks, and lunches on 27/28/29 February 2012 as well as the social programme on Wednesday, 29 February 2012. The participation in the following conference dinner is optional and requires extra registration (additional costs per person: 48 Euro; for more information please see the paragraph „Social programme and conference dinner“ on the next page).

#### Registration category

#### Late & on-site registration fee (from 27 January 2012)

	incl. 19 % VAT	excl. VAT*
Late & on-site registration for members	530 €	445.38 €
Late & on-site registration for non-members	610 €	512.61 €
Late & on-site registration for student members	290 €	243.70 €
Late & on-site registration for student non-members	310 €	260.50 €
Late & on-site registration for one-day	280 €	235.29 €

**\* PLEASE NOTE:** Registrations from companies and non-university research institutes registered in EU countries (except Germany) are exempted from VAT, if **VAT no. is given**.

### ORAL PRESENTATIONS

**Time slots:** Presenting authors are allotted 20 minutes (15 minutes presentation plus 5 minutes for discussion). Please plan your presentation accordingly to meet the 20 minutes maximum.

**Presentation upload:** Speakers are requested to upload their presentation to the computer in the meeting room well in advance to their talk.

**Presentation format:** Please bring your presentation on an USB mass storage, CD-ROM or DVD and include all video files. File formats: ppt, pptx and pdf. A Windows-based presentation computer will be provided.

**For Mac users:** To make sure your presentation is displayed correctly, please:

- bring your presentation as pdf-file with fonts embedded or
- restrict yourself to Arial/Times New Roman (not Times)/Courier New (not Courier)/Symbol/Windings when creating your ppt- or pptx-file.

**Technical equipment:** All technical equipment (presentation computer, video projector, sound system, laser pointer) will be available on-site. It is not possible to use your personal laptop.

### POSTER PRESENTATIONS

Poster authors are requested to be present at their posters during the official poster session. Poster set-up and removal is in the responsibility of the authors. The poster session will be taking place on Tuesday, 28 February, 18:00 - 19:30. Poster strips will be provided by the organiser.

**Required poster size:** The posters should have a size of DIN A1 (width: 59,4 cm x height: 84,1 cm) or DIN A0 (width: 84,1 cm x height: 118,9 cm) in a portrait format. The size of the poster boards is 94 cm x 193 cm (width x height).


### EOS TOPICAL MEETING DIGEST

The registration includes a CD-ROM with the abstracts of all accepted, invited and plenary presentations at this topical meeting (ISBN 978-3-00-033711-6).

NOTE: A one-day registration does not include the digest CD-ROM. It can be ordered separately.

The EOS does not publish proceedings with extensive papers. Authors wishing to publish in-depth papers are welcome to take advantage of the special publication offer for JEOS:RP (see paragraph "Special publication fee for attendees"). This publication offer is an option but no obligation.

### JEOS:RP - SPECIAL SPECIAL PUBLICATION OFFER



All attendees of DO 2012 will receive a **20% discount** on the publication rate for the e-journal of the European Optical Society: Rapid Publications (JEOS:RP, [www.jeos.org](http://www.jeos.org)). The paper submitted to JEOS:RP must be an original contribution that is connected to one of the conference topics and must be submitted no later than 2 May 2012.

#### Special publication fee for attendees

280 € (instead of 350 €) for full EOS members

320 € (instead of 400 €) for non EOS members

### BEST STUDENT PRESENTATION AWARD



The two best student presentations (oral and poster) at this EOS Topical Meeting will be awarded a diploma, an EOS student membership for 2012 and a prize sponsored by the publisher Springer. All student presenters are eligible to the prize. The winners will be announced at the end of the last session on Thursday.

### WIFI ACCESS

Free WIFI access will be available in the Aula. **Network name:** TUDelft-congress | **Access code:** jpesk.

### SOCIAL PROGRAMME & CONFERENCE DINNER



The social programme on **Wednesday, 29 February 2012, from 17.00 -18.30** includes a guided tour of the Royal Delft, Koninklijke Porceleyne Fles, the last remaining Delftware factory from the 17th century. The world famous Royal Delftware is still entirely hand-painted according to centuries-old tradition and all participants will attend the production process with a painting demonstration.

The meeting point for the social programme is in the foyer of the Conference Centre at 16:45.

**Location:** Rotterdamseweg 196

The conference dinner will be taking place in the Partycentrum - Restaurant de Brasserie on **Wednesday, 29 February, starting at 18:30** with welcome drinks and appetizers.

**Participation in the conference dinner requires extra registration** (deadline: 17 February 2012). Costs per person: 48 Euro. The fee includes an international style buffet with cold & warm food as well as beverages. Spirits must be paid separately.

**Location:** Brasserskade 2a

To register please use the fax form at [www.myeos.org/events/do2012](http://www.myeos.org/events/do2012) or register through the EOS Online Shop at [www.myeos.org/shop](http://www.myeos.org/shop).



### WELCOME RECEPTION



All attendees of DO 2012 are cordially invited to attend the welcome reception - sponsored by TNO - on **Monday, 27 February 2012, from 18.00 - 20.00**.

**Location:** Atrium at TNO (adjacent building to the TU Delft)  
Stieltjesweg 1

WHISKEY TASTING



EOSAM 2012 moves to Aberdeen

In 2012 the EOS Annual Meeting moves from Paris to Aberdeen. For this reason, the office will offer a whisky tasting for all attendees of this topical meeting.

You will find it during the last coffee break on **Wednesday, 29 February, 15:50 - 16:30**.

EOS INTERNAL MEETING SCHEDULE

- EXECOM MEETING
 

	Location: Conference Centre, Delft University of Technology Netherlands
Thursday, 1 March	Room: Commissiekamer 1.
14:00-17:00	
  
- BOARD MEETING
 

	Location: Conference Centre, Delft University of Technology Netherlands
Friday, 2 March	Room: Commissiekamer 3.
11:00-15:00	



Nedinsco develops, designs and builds sophisticated, optical, mechanical and electronic systems (including software) for the global defence, diagnostics and semiconductor markets.

- **Founded in 1921**
- **Development & design**
- **Proto & serial production**
- **High precision assembly**
- **Qualification & test**

Nedinsco has an opportunity available for those looking for personal development and exciting challenges:

**OPTICAL ENGINEER**



Recognized for our continued focus on innovation, we offer ambitious individuals the perfect platform to develop their skills and contribute in an open and learning environment.

For further details on the above position and to apply please visit our website, use the QR code above or contact our Human Resources department via e-mail: [hr@nedinsco.nl](mailto:hr@nedinsco.nl)

**nedinsco.com**

**photonic technology**

## WELCOME TO DO 2012

After 20 years of active research and development of design and fabrication techniques, diffractive optics is approaching the status of a mature technology. In addition to the ongoing fundamental basic and applied research, we can find diffractive optics in increasingly many applications which are either completely based on diffraction phenomena or for which the diffraction solution is remarkably more compact, light-weight, or cheaper than its alternatives.

This meeting will be the eighth in a series. Previous meetings took place in Savonlinna (1997), Jena (1999), Budapest (2001), Oxford (2003), Warsaw (2005), Barcelona (2007) and Koli (2010).

The 2012 meeting in Delft includes a focus session on Diffraction Optics in the High-Tech Industry. This session gathers scientists and engineers working in the field of diffractive optics in industry and those in academy that work on industrial problems. The Topical Meeting will give attendees a wide picture of Diffractive Optics, from fundamental modelling of light-matter interaction to design techniques and from fabrication techniques to novel applications.

### FOCUS SESSION

Diffractive Optics in the High-Tech Industry

#### TOPICS

- Novel optical materials including metamaterials
- Modelling of diffractive optics
- Optical beam shaping
- Polarization
- Scattering
- Diffractive Optics for Solid State Lighting and solar Photonic crystals
- Adaptive and switchable diffractive optics
- Diffractive Optics for Life Sciences and sensing
- Inverse problems in diffraction optics
- Fabrication and characterization
- Measurements and inspection

#### GENERAL CHAIRS



**Paul Urbach**  
Chair of Diffractive Optics 2012  
Delft University of Technology (NL)



**Jani Tervo**  
Chair of Diffractive Optics 2012  
University of Joensuu (FI)

#### PROGRAMME COMMITTEE

- **Juan Campos Coloma**, Universidad Autònoma de Barcelona (ES)
- **Pierre Chavel**, Centre National de la Recherche Scientifique (FR)
- **Zbigniew Jaroszewicz**, Institute of Applied Optics (PL)
- **Lifeng Li**, Tsingua University (CN)
- **Masud Mansuripur**, University of Arizona (USA)
- **Silvania Pereira**, Delft University of Technology (NL)
- **Jari Turunen**, University of Joensuu (FI)
- **Frank Wyrowski**, University of Jena (DE)



## PLENARY SPEAKER

Monday, 27 February 2012, Senaatszaal

08:35 - 09:20



Martin Wegener, Karlsruhe Institute of Technology (DE)

**3D photonic metamaterials by diffraction-unlimited laser lithography**

We review our recent efforts towards diffraction-unlimited far-field optical lithography based on stimulated-emission-depletion (STED) inspired direct laser writing (DLW). This has, for example, enabled the first three-dimensional polarization-independent visible-frequency invisibility cloak (for the light amplitude and phase).

## INVITED SPEAKERS

Monday, 27 February 2012, Senaatszaal

14:00 - 14:30



Jakob Stamnes, University of Bergen (NO)

**Wave propagation, diffraction, and focusing in uniaxially and biaxially anisotropic media**

In many applications, such as wave plates, resonators, electro-optic modulators, polarizers, and thin-film couplers, it is important to have an accurate description of the propagation of electromagnetic beams in anisotropic media. Therefore, considerable attention has been devoted to these issues in the past.

Tuesday, 28 February 2012, Senaatszaal

08:30 - 09:00



Allard Mosk, University of Twente (NL)

**Scattering Lenses**

Scattering of light by random collections of particles or surface roughness can be used to increase the resolution of imaging systems. We demonstrate 100-nm resolution focusing of light by scattering in a high-index material.

14:00 - 14:30



Bernd Kleemann, Carl Zeiss AG (DE)

**Perfect blazing for echelle gratings in Littrow mount**

Perfect blazing for echelle gratings in Littrow mount really exists, even simultaneously in TE and TM polarization. Then no directed stray light from other diffraction orders occurs. In the ideal case of infinite conductivity, 100% efficiency is diffracted into the Littrow order. For real metal gratings, a small loss occurs.

Wednesday, 29 February 2012, Senaatszaal

08:30 - 09:00



Cesare Umeton, Università della Calabria (IT)

**All optical switchable diffraction gratings realized in liquid crystalline composite structure with metamaterial applications**

We report on the fabrication and all optical characterization of two different kinds of switchable diffraction gratings containing light responsive Liquid Crystals. We foresee that, when doped with gold nanoparticles, these structures can be exploited to obtain the tuning of "active" plasmonic responses, to be used for metamaterial applications.

INVITED SPEAKERS

Wednesday, 29 February 2012, Senaatszaal

14:00 - 14:30



**Robert Magnusson**, University of Texas in Arlington (US)

**Theory, experiment, and applications of photonic resonance effects in nanopatterned films-A review**

Fundamental photonic resonance effects in thin periodic films with nanoscale features are presented and examples of associated spectral expressions given. Potential applications including biosensors and optical components are discussed.

Thursday, 1 March 2012, Senaatszaal

08:30 - 09:00



**Makoto Kuwata-Gonokami**, University of Tokyo (JP)

**Polarization control with artificial chiral structures**

Artificial sub-wavelength two dimensional grating structures without mirror symmetry exhibit strong optical activity and rotate polarization states of propagating optical and terahertz waves. We also demonstrate circularly polarized spontaneous emission of quantum dots embedded in a semiconductor chiral photonic crystal.

10:30 - 11:00



**Jaime Gomez-Rivas**, FOM Institute AMOLF (NL)

**Modified quantum dot emission with arrays of plasmonic nanoparticles**

We present an experimental study of the modified emission of a layer of quantum dots (QDs), just a few nanometers thick, with plasmonic structures. The coupling of localized surface plasmon resonances in metallic particles through in-plane diffraction in arrays of these particles leads to the formation of quasi-bound surface states known as surface lattice resonances (SLRs). The recent observation of extremely narrow SLRs in arrays of gold nanoparticles has created expectations for these arrays as plasmonics sensors. We show here that these resonances can lead to a strong modification of the emission due to the modified density of optical states at which the excited QDs can decay and the diffractive coupling of SLRs into the far-field.

Publish your research with JEOS:RP

New impact factor  
2010: 1.044



JOURNAL OF  
THE EUROPEAN  
OPTICAL SOCIETY  
RAPID PUBLICATIONS

Discounted publication rates for  
attendees of DO 2012

Special publication rates:  
(incl. 20 % discount)

- 320 € (non-member rate)
- 280 € (member rate)

The paper submitted must be an original contribution  
that is connected to the topics of this EOS event.

Paper submission deadline:  
2 May 2012

Journal Management Contact: Silke Kramprich  
Phone: +49-511-2788-117 | Email: jeos-rp@myeos.org

[www.jeos.org](http://www.jeos.org)

**SUNDAY, 26 FEBRUARY 2012**

16:00 - 18:00 Pre-Registration

**MONDAY, 27 FEBRUARY 2012**

08:30 - 08:35 Opening by the Chairs

08:35 - 09:20 Plenary Talk

**3D photonic metamaterials by diffraction-unlimited laser lithography**

Martin Wegener, Karlsruhe Institute of Technology (DE)

09:20 - 10:20 Session I

10:20 - 10:50 Coffee break

10:50 - 12:10 Session II

12:10 - 14:00 Lunch break

14:00 - 15:50 Session III

15:50 - 16:20 Coffee break

16:20 - 18:00 Session IV

18:00 - 20:00 Welcome Reception sponsored by TNO

**TUESDAY, 28 FEBRUARY 2012**

08:30 - 09:40 Session V

09:40 - 10:10 Coffee break

10:10 - 11:50 Session VI

11:50 - 13:40 Lunch break

13:40 - 15:10 Session VII

15:10 - 16:00 Coffee break

16:00 - 17:40 Session VIII

18:00 - 19:30 Poster session

**WEDNESDAY, 29 FEBRUARY 2012**

08:30 - 10:00 Session IX

10:00 - 10:30 Coffee break

10:30 - 12:10 Session X

12:10 - 14:00 Lunch break

14:00 - 15:50 Session XI

15:50 - 16:30 Coffee break &amp; Whiskey tasting

17:00 - 18:30 Social Programme

18:30 - 22:30 Conference Dinner

**THURSDAY, 1 MARCH 2012**

08:30 - 10:00 Session XII

10:00 - 10:30 Coffee break

10:30 - 12:20 Session XIII

12:50 END OF EOS TOPICAL MEETING

Senaatszaal, TU Delft

08:30 - 08:35 OPENING BY THE CHAIRS

Jani Tervo, University of Eastern Finland (FI)  
Paul Urbach, Delft University of Technology (NL)

NOTES

08:35 - 09:20

PLENARY TALK

3D photonic metamaterials by diffraction-unlimited laser lithography

*J. Fischer, T. Ergin, M. Wegener; Institut für Angewandte Physik, Institut für Nanotechnologie, and DFG-Center for Functional Nanostructures (CFN) Karlsruhe Institute of Technology (KIT) (DE).*  
We review our recent efforts towards diffraction-unlimited far-field optical lithography based on stimulated-emission-depletion (STED) inspired direct laser writing (DLW). This has, for example, enabled the first three-dimensional polarization-independent visible-frequency invisibility cloak (for the light amplitude and phase). [4881]

09:20 - 10:20

SESSION I

Chair: Paul Urbach, Delft University of Technology (NL)

09:20 - 09:40

STUDENT PRESENTATION

Time-domain measurement of diffraction phenomena

*P. Piksarv<sup>1</sup>, M. Lähmus<sup>1</sup>, H. Valtma-Lukner<sup>1</sup>, B. Bowlan<sup>2</sup>, R. Trebino<sup>2</sup>, P. Saari<sup>1,3</sup>; <sup>1</sup>Institute of Physics, University of Tartu (EE); <sup>2</sup>School of Physics, Georgia Institute of Technology (US); <sup>3</sup>Estonian Academy of Sciences (EE).*

Employing the SEA TADPOLE measurement technique, we directly measure the spatio-temporal field of ultrashort pulses after diffracting off simple apertures and pulsed Bessel beams generated by circularly symmetric diffraction gratings. Our measurements beautifully confirm the time-domain treatment of diffraction. [4871]

09:40 - 10:00

Investigating Phase Talbot Image of Wavelength-Scale Period Amplitude Grating

*M.-S. Kim<sup>1</sup>, T. Scharf<sup>1</sup>, H.P. Herzig<sup>1</sup>, C. Menzel<sup>2</sup>, C. Rockstuhl<sup>2</sup>; <sup>1</sup>Ecole Polytechnique Fédérale de Lausanne (EPFL), Optics & Photonics Technology Laboratory (CH); <sup>2</sup>Friedrich-Schiller-Universität Jena, Institute of Condensed Matter Theory and Solid State Optics (DE).*

We interferometrically measure the Talbot length of amplitude gratings with periods comparable to the wavelength. By recording phase images in real-space with superresolution, the Talbot length can be determined with an unprecedented precision. Rigorous simulations verify all experiments. The measurements allow us to access the predictive strength of analytical expressions for the Talbot lengths and show that these expressions maintain their accuracy well beyond the limits usually anticipated. [4827]

10:00 - 10:20

Stokes imaging polarimetry with a single-pixel detector

*V. Durán<sup>1,2</sup>, P. Clemente<sup>1,3</sup>, M. Fernández<sup>1,2</sup>, E. Tajahuerce<sup>1,2</sup>, J. Lancis<sup>1,2</sup>, P. Andrés<sup>4</sup>; <sup>1</sup>Institut de Noves Tecnologies de la Imatge (INIT), Universitat Jaume I (ES); <sup>2</sup>GROC - UJI, Departament de Física, Universitat Jaume I (ES); <sup>3</sup>Servei Central d'Instrumentació Científica, Universitat Jaume I (ES); <sup>4</sup>Departamento de Óptica, Universitat de València (ES).*

We present an optical system that performs spatially resolved measurements of the Stokes parameters of light with a single-pixel detector. This fact is possible by applying the theory of compressive sampling. Experimental results are presented for an object that provides an inhomogeneous polarization distribution. [4903]

10:20 - 10:50 Coffee break

10:50 - 12:10

SESSION II

Chair: Lifeng Li, Tsinghua University (CN)

10:50 - 11:10

STUDENT PRESENTATION

Design of a circular wire grid grating to generate radially polarized light in the near UV region

*K. Ushakova, S.F. Pereira, H.P. Urbach; Delft University of Technology, Faculty of Applied Sciences, Department of Imaging Science & Technology, Optics Research Group (NL).*

Formation of a high quality radially polarized light in the near UV wavelength region (405 nm, 365 nm, 193 nm) using thin metal film diaphragm compound of sub wavelength concentric nanoslit grooves is inspected by using subsequently models of ideal metal-insulator-metal waveguide, 1D grating and 3D diaphragm configuration. [4862]



## Senaatszaal, TU Delft

11:10 - 11:30

**Phase retardation in diffracted light**

K. Ventola<sup>1</sup>, J. Tervo<sup>1</sup>, S. Siitonen<sup>2</sup>, M. Kuittinen<sup>1</sup>; <sup>1</sup>University of Eastern Finland, Dep. of Physics and Mathematics (FI); <sup>2</sup>Nanocomp Ltd (FI).

Phase retardation occurring in non-zeroth diffraction orders of 3-D dielectric gratings is studied numerically and experimentally. An element producing half-waveplate effect in -2nd reflected order is presented. [4879]

11:30 - 11:50

**Security diffractive elements – Nanogravure of 2nd kind**

L. Kotačka, P. Vizdal, T. Běhounek; Optaglio s.r.o. (CZ).

The paper summarizes recent development on the synthesized holographic and diffractive security elements. Such security devices are recorded by means of the electron beam lithography with the resolution 2,500.000 dpi, thus with particular resolution of the beam spot 10 nm. This allows to originate very unique devices serving for anti-counterfeit measures, where some special and/or rudimentary diffractive features are synthetically introduced to other security diffractive elements. Such security holograms are reflecting the highest requirements for protection of governmental documents and banknotes. [4880]

11:50 - 12:10

**Improved resolution for soft x-ray spectroscopy: single-order operation of lamellar multilayer gratings**

R. van der Meer<sup>1</sup>, I.V. Kozhevnikov<sup>2</sup>, B. Krishnan<sup>1</sup>, J. Huskens<sup>1</sup>, M.J. de Boer<sup>1</sup>, B. Vratzov<sup>3</sup>, P.E. Hegeman<sup>4</sup>, G.C.S. Brons<sup>4</sup>, H.M.J. Bastiaens<sup>1</sup>, K.-J. Boller<sup>1</sup>, F. Bijkerk<sup>1,5</sup>; <sup>1</sup>MESA+ Institute for Nanotechnology, University of Twente (NL); <sup>2</sup>Institute of Crystallography, Russian Academy of Sciences (RU); <sup>3</sup>NT&D - Nanotechnology and Devices (DE); <sup>4</sup>PANalytical (NL); <sup>5</sup>FOM-Institute Rijnhuizen (NL).

Spectral resolution improvements of nearly a factor four using Lamellar Multilayer Gratings in the soft x-ray range are reported. The design is based on single-order operation, in which the incident beam effectively only excites a single diffraction order. Measured bandwidths are in good agreement with the theoretical model. [4878]

12:10 - 14:00 Lunch break

14:00 - 15:50

**SESSION III**

Chair: Martin Wegener, Karlsruhe Institute of Technology (DE)

14:00 - 14:30

Invited talk

**Wave propagation, diffraction, and focusing in uniaxially and biaxially anisotropic media**

J.J. Stamnes; Department of Physics and Technology, University of Bergen (NO).

In many applications, such as wave plates, resonators, electro-optic modulators, polarizers, and thin-film couplers, it is important to have an accurate description of the propagation of electromagnetic beams in anisotropic media. Therefore, considerable attention has been devoted to these issues in the past. [4883]

14:30 - 14:50

**Fourier-space filtering in classical ghost imaging and diffraction**

T. Shirai<sup>1</sup>, H. Kellock<sup>2</sup>, T. Setälä<sup>2</sup>, A.T. Friberg<sup>2,3,4</sup>; <sup>1</sup>Electronics and Photonics Research Institute, National Institute of Advanced Industrial Science and Technology (AIST) (JP); <sup>2</sup>Department of Applied Physics, Aalto University (FI); <sup>3</sup>Department of Physics and Mathematics, University of Eastern Finland (FI); <sup>4</sup>Department of Microelectronics and Applied Physics, Royal Institute of Technology (KTH) (SE).

We show that introduction of a spatial filter into the reference arm of the conventional classical ghost-imaging setup allows for imaging of phase objects, while a spatial filter in the conventional classical ghost-diffraction arrangement enables imaging through phase aberrations, whether deterministic or random. [4950]

14:50 - 15:10

**A generalization of the van Cittert-Zernike theorem and the interconnection between the temporal- and spatial longitudinaltransverse degrees of partial coherence**

B.J. Hoenders<sup>1</sup>, Y. Cai<sup>2</sup>, Y. Dong<sup>2</sup>; <sup>1</sup>University of Groningen, Center for Theoretical Physics and Zernike Institute for Advanced Materials (NL); <sup>2</sup>School of Physical Sciences and Technology, Soochow University (CN).

A generalization of the Zernike-van Cittert theorem is used to analyse the interdependence between the temporal- longitudinal- and transverse spatial coherence. We use a model which assumes that the mutual coherence function is spatial incoherent at a planar surface and that the temporal coherence is not only quasi monochromatic (as in the Zernike-van Cittert case), but contains temporal correlation terms as well. [4907]

NOTES

## Senaatszaal, TU Delft

15:10 - 15:30

**Spatial correlation for propagated fields and the phase problem**

*O. El Gawhary<sup>1,2</sup>, A. Wiegmann<sup>3</sup>, N. Kumar<sup>1</sup>, S.F. Pereira<sup>1</sup>, H.P. Urbach<sup>1</sup>; <sup>1</sup>Optics Research Group, Delft University of Technology (NL); <sup>2</sup>Dutch Metrology Institute (VSL) (NL); <sup>3</sup>Physikalisch-Technische Bundesanstalt (PTB) (DE).*

Measuring the phase of an optical field it is known to be not an easy task. In fact, any common detector shows integration times much longer than the typical period of an electromagnetic field in the visible. Although there are methods that allow to turn the phase information into intensity one, such as interferometry, often specific applications require solutions easier from an implementation point of view. In optics, retrieving the phase of a field has important implications in microscopy as well as in the analysis of the performances of an optical system, as, for instance, in the measurement of its aberrations. [4956]

15:30 - 15:50

**Resist based aberration metrology using Phase-Shift Gratings**

*S. van Haver<sup>1</sup>, W. Coene<sup>1</sup>, P. van Adrichem<sup>2</sup>, L. de Winter<sup>3</sup>; <sup>1</sup>ASML, TE RES Research SMC (NL); <sup>2</sup>ASML, TE SE Focus (NL); <sup>3</sup>ASML, DE EUV ISQ Reticles&Imaging (NL).*

We propose a method to reconstruct the aberrations of a lithographic system through wafer overlay metrology. We designed new 1D and 2D phase-shift gratings (PSG's) that translate a phase difference of two diffracted beams into an overlay error of the resulting resist pattern. The combined information of the lateral image position for many of our new 1D and 2D PSG's allows for reconstruction of the aberration function of the optical imaging system of the scanner. [4947]

15:50 - 16:20 Coffee break

16:20 - 18:00

**SESSION IV**

*Chair: Jani Tervo, University of Joensuu (FI)*

16:20 - 16:40

**High-NA diffractive array illuminators**

*S. Stallinga; Quantitative Imaging Group, Department of Imaging Science and Technology, Delft University of Technology (NL).*

Binary phase diffractive array illuminators are studied, which can produce large area arrays of high-NA spots. These array illuminators may serve as key component in costeffective imaging systems. [4932]

16:40 - 17:00

**Planar Diffractive Diffuser**

*V. Zagolla, C. Moser; EPFL, LAPD (CH).*

We aim to investigate the use of a novel Red, Green, Blue sensitive holographic photopolymer for creating diffracting structures providing diffuse illumination and light trapping over the whole visible spectrum, targeting an increase of light absorption in polymer/dye-sensitized solar cells. [4860]

STUDENT PRESENTATION

17:00 - 17:20

**Diffractive optics for lighting applications**

*H.J. Cornelissen<sup>1</sup>, C. Ho<sup>2</sup>, N. Haj-Hosseini<sup>2</sup>; <sup>1</sup>Philips Research, Optics Department (NL); <sup>2</sup>Delft University, Optics Research (NL).*

Sub-micron diffraction gratings have been used for two illumination application. One is to create a transparent see-through luminaire which can be used to illuminate and read a paper document. A second is a light sensor that can be used in a feedback loop to control a multicolor LED lamp. Optical design and prototypes are presented. [4976]

17:20 - 17:40

**Optical design with smooth and diffractive optical surfaces by generalized ray tracing**

*M.A. Golub; Tel Aviv University, Department of Electrical Engineering (IL).*

A generalized ray tracing method based on calculation of local wavefront curvatures was developed. Sequentially cascaded curved refractive optical surfaces with nonsymmetrical diffractive phase functions can now be designed with nearly the same simplicity as customary in paraxial optics of thin lenses and Fourier optics. [4940]

NOTES

## Senaatszaal, TU Delft

17:40 - 18:00

STUDENT PRESENTATION

**Exact simulation of the diffraction of large, high-NA DOE-sections with  $O(N \log N)$  time and memory resort**

*A.V. Tishchenko<sup>1</sup>, A.A. Shcherbakov<sup>2</sup>; <sup>1</sup>University of Lyon, Lab. H. Curien UMR CNRS 5516 (FR); <sup>2</sup>Moscow Institute of Physics and Technology (RU).*

A major jump in the capacity of exactly solving 3D diffraction problems in microoptics is achieved by a unique chain of mathematical steps permitting to avoid any matrix inversion. This powerful new method exhibiting a linear CPU time dependence on the number of considered diffraction orders enables the instant simulation of 2D high-NA DOE-sections large enough for an exact and meaningful representation/optimization of the optical function of the complete element under arbitrary incidence conditions. [4858]

NOTES

18:00 - 20:00 Welcome reception - sponsored by TNO



**Venue:** Atrium at TNO  
(adjacent building to the TU Delft),  
Stieltjesweg 1

NOTES

Senaatszaal, TU Delft

08:30 - 09:40

SESSION V

Chair: Ari T. Friberg, University of Eastern Finland, Aalto University (FI) & Royal Institute of Technology (KTH) (SE)

08:30 - 09:00

Invited talk

Scattering Lenses

E.G. van Putten<sup>1</sup>, D. Akbulut<sup>1</sup>, J. Bertolotti<sup>1,2</sup>, W.L. Vos<sup>1</sup>, A. Lagendijk<sup>1,3</sup>, A.P. Mosk<sup>1</sup>; <sup>1</sup>Complex Photonic Systems, Faculty of Science and Technology and MESA+ Institute for Nanotechnology, University of Twente (NL); <sup>2</sup>Dipartimento di Fisica, University of Florence (IT); <sup>3</sup>FOM Institute for Atomic and Molecular Physics (AMOLF) (NL).

Scattering of light by random collections of particles or surface roughness can be used to increase the resolution of imaging systems. We demonstrate 100-nm resolution focusing of light by scattering in a high-index material. [4875]

09:00 - 09:20

Compensation of spatio-temporal light spreading in second harmonic generation with diffractive optical elements

R. Martínez-Cuenca<sup>1</sup>, J. Pérez-Vizcaino<sup>1</sup>, O. Mendoza-Yero<sup>1</sup>, B. Alonso<sup>2</sup>, Í.J. Sola<sup>2</sup>, G. Mínguez-Vega<sup>1</sup>, J. Lancis<sup>1</sup>, P. Andrés<sup>3</sup>; <sup>1</sup>GROC\*UJI, Institut de Noves Tecnologies de la Imatge (ES); <sup>2</sup>Dept. de Física Aplicada, Universidad de Salamanca (ES); <sup>3</sup>Dept. de Òptica, Universitat de València (ES).

We propose a technique for one-shot generation of arbitrary second harmonic (SH) patterns based on diffractive optical elements (DOEs) and ultrafast pulses. We show that for pulse durations fewer than 100 fs spatio-temporal shaping is mandatory to overcome dispersive effects that spread irradiance patterns both in space and time. [4904]

09:20 - 09:40

STUDENT PRESENTATION

Glass Metal Nanocomposites Based Diffraction Gratings

M.I. Petrov<sup>1</sup>, Y.P. Svirko<sup>1</sup>, A.A. Lipovskii<sup>2</sup>, O.V. Shustova<sup>2</sup>; <sup>1</sup>University of Eastern Finland (FI); <sup>2</sup>St.-Petersburg State Polytechnical University (RU).

In this work we investigate the implementation of glasses with embedded metal nanoparticles for diffraction gratings fabrication. Spectral properties of diffraction strongly depend on nanocomposite parameters such as filling factor and nanoparticle diameter. Fabrication technics of nanocomposites based gratings are also discussed. [4938]

09:40 - 10:10 Coffee break

10:10 - 11:50

SESSION VI

Chair: Zbigniew Jaroszewicz, Institute of Applied Optics (PL)

10:10 - 10:30

Diffractive optical elements for testing cylindrical surfaces

A.G. Poleshchuk<sup>1</sup>, R.K. Nasyrov<sup>1</sup>, J.-M. Asfour<sup>2</sup>; <sup>1</sup>Institute of Automation & Electrometry SB RAS (RU); <sup>2</sup>Dioptric GmbH (DE).

Different types of DOEs and optical test layouts for testing cylinder surfaces have been developed, investigated and experimentally approved. Results for design and fabrication of holograms for cylindrical mirrors testing are presented. [4916]

10:30 - 10:50

Towards absolute interferometry for the testing of freeform surfaces

C. Pruss, S. Peterhänsel, F. Schaal, W. Osten; Universität Stuttgart, Institut für Technische Optik (DE).

One of the most accurate methods to test steep high-precision freeform surfaces in an interferometer is the well established null test using computer generated holograms (CGH). In this contribution we discuss error sources and ways for their separation from the errors of the surface under test. [4945]

10:50 - 11:10

STUDENT PRESENTATION

Semiconductor metrology with inverse diffraction approach "Coherent Fourier Scatterometry"

N. Kumar<sup>1</sup>, O.El. Gawhary<sup>1</sup>, S.F. Pereira<sup>1</sup>, W. Coene<sup>2</sup>, H.P. Urbach<sup>1</sup>; <sup>1</sup>Delft University of Technology (NL); <sup>2</sup>A.S.M.L BV (NL).

A new approach for optical far field analysis and its implementation is presented here. Under suitable conditions, using focused beam from a spatially coherent light source can increase the accuracy of the optical Scatterometry technique. [4900]

NOTES

## Senaatszaal, TU Delft

11:10 - 11:30

**Analysis of Scanning and Polarization Effects in Coherent Fourier Scatterometry**

*S. Roy, O. El Gawhary, S.F. Pereira, H.P. Urbach; Optics Research Group, Department of Imaging Science and Technology, Delft Institute of Technology (NL).*

Coherent Fourier Scatterometry (CFS) is a promising candidate for high accuracy nano-metrology as this method offers accuracy, convenience of measurement and fast processing time simultaneously. In this paper we show that the use of coherence and multiple scanning makes CFS more sensitive than incoherent scatterometry and how an optimum number of scanning positions can be predicted for a specific case. The sensitivity of the method can be further increased by making the incident and measured fields cross-polarized. [4893]

11:30 - 11:50

STUDENT PRESENTATION

**Hyperchromatic Confocal Sensor Systems**

*M. Hillenbrand, C. Wenzel, X. Ma, P. Feßler, S. Sinzinger; TU Ilmenau, IMN MacroNano® (DE).*

We discuss the potential and challenges of diffractive and hybrid systems for hyperchromatic sensor applications. Design, fabrication, and characterization of chromatic confocal sensors for innovative distance and wall thickness measurement systems are presented. [4847]

11:50 - 13:40 Lunch break

13:40 - 15:10

## SESSION VII

*Chair: Benfeng Bai, Tsinghua University (CN) & University of Eastern Finland (FI)*

13:40 - 14:10

Invited talk

**Perfect blazing for echelle gratings in Littrow mount**

*B.H. Kleemann; Carl Zeiss AG, Corporate Research and Technology (DE).*

Perfect blazing for echelle gratings in Littrow mount really exists, even simultaneously in TE and TM polarization. Then no directed stray light from other diffraction orders occurs. In the ideal case of infinite conductivity, 100% efficiency is diffracted into the Littrow order. For real metal gratings, a small loss occurs. [4835]

14:10 - 14:30

**Electromagnetic edge singularity and Rayleigh expansion**

*L. Li; Tsinghua University, Department of Precision Instruments (CN).*

When irregular electric field singularity at the edge of a wedge composed of lossless metal and dielectric media is excited, all grating modeling methods that use Rayleigh expansion do not converge. This work generalizes a recent work to cases of an arbitrary wedge angle. Apparent conflict with Rayleigh expansion is also discussed. [4922]

14:30 - 14:50

**Recent progress in numerical modeling of gratings. Should we abandon the MMFE?**

*G. Granet; Clermont Université, Université Blaise Pascal, Lasmea (FR) & Cnrs, UMR 6602, Lasmea (FR).*

We report on significant improvements of various numerical methods for diffraction gratings. [4914]

14:50 - 15:10

STUDENT PRESENTATION

**RMCA: an alternative algorithm for the fast computation of 2D gratings**

*B. Portier<sup>1</sup>, F. Pardo<sup>2</sup>, Riad Haïdar<sup>1,3</sup>, J.-L. Pelouard<sup>2</sup>; <sup>1</sup>ONERA – The French Aerospace Lab (FR); <sup>2</sup>Laboratoire de Photonique et de Nanostructures (LPN-CNRS) (FR); <sup>3</sup>Ecole Polytechnique (FR).*

Calculation speed is often an issue when simulating optical properties of complex nanostructures, such as 2D lamellar gratings. Here, we present a new modal method which enables fast computation of these properties without loss of precision. The Rigorous Maxwell with Constitutive Approximation (RMCA) is based on the exact discrete formulation of the Maxwell equations, in which we introduce polynomial approximations of the constitutive equations to obtain the eigenproblem equation in each layer. The use of sparse matrices enables us to significantly enhance the calculation speed by calculating only a short number of eigenmodes, and numerical results on published examples show that we can have the same precision as with the RCWA method in less time. [4928]

15:00 - 16:00 Coffee break

NOTES

Senaatszaal, TU Delft

16:00 - 17:40

SESSION VIII

Chair: Pierre Chavel, Centre National de la Recherche Scientifique (FR)

16:00 - 16:20

**Molding of Diffractive Glass Lenses**

D. Hollstegge, B. Bulla, M. Hüntel, O. Dambon, F. Klocke; Fraunhofer Institute for Production Technology (DE).

The trend towards miniaturization and functional integration of optical systems and components is also reflected in the increased demand for structured glass optics. Precision glass molding can be used to produce full glass diffractive optics, especially hybrid lenses such as asphero-diffractive lenses. [4944]

16:20 - 16:40

STUDENT PRESENTATION

**Nearly temperature independent waveguide gratings**

M.R. Saleem<sup>1,3</sup>, M.B. Khan<sup>3</sup>, S. Honkanen<sup>1,2</sup>, J. Turunen<sup>1</sup>; <sup>1</sup>University of Eastern Finland, Department of Physics and Mathematics (FI); <sup>2</sup>Aalto University, Department of Micro and Nanosciences (FI); <sup>3</sup>National University of Science and Technology, School of Chemical and Materials Engineering (PK).

We investigated nearly athermalized organic-inorganic devices as athermal guided mode resonance filters GMRFs. The spectral reflectance efficiency is observed to be nearly independent on temperature over tens of degrees above room temperature. Two important factors are considered for the shift in the resonance peak; mainly thermal expansion TEC and thermo-optic TOC coefficients of the selected materials. [4887]

16:40 - 17:00

**Advanced matrix laser lithography for fabrication of photonic micro-structures**

M. Škereň, J. Svoboda, P. Fiala; Czech Technical University in Prague, Faculty of Nuclear Sciences and Physical Engineering, Department of Physical Electronics (CZ).

The laser lithography technique is presented, which enables to fabricate large area periodic and aperiodic micro-structures for various photonic applications. The device based on imaging the micro-structure from a computer driven spatial light modulator is presented together with experimental demonstration of fabricated structures. [4925]

17:00 - 17:20

**Self-organisation of Nematic Gold Nanoparticles in Periodic Polymeric Structures**

R. Caputo<sup>1</sup>, B.J. Tang<sup>2</sup>, L. De Sio<sup>1</sup>, G.H. Mehl<sup>2</sup>, C. Umeton<sup>1</sup>; <sup>1</sup>University of Calabria, LICRYL (Liquid Crystals Laboratory, IPCF-CNR), Center of Excellence (CEMIF.CAL) and Department of Physics (IT); <sup>2</sup>University of Hull, Department of Chemistry (GB).

We report on the self-organisation of Nematic Gold nanoparticles in host liquid crystal phases. Experiments show that these particles have a very good miscibility with several liquid crystal phases (nematics, cholesterics and smectics). Periodic channelled structures allowed checking the liquid crystalline behaviour of obtained materials. [4911]

17:20 - 17:40

**Stitchingless grating at the wall of a cylinder by single-shot radial phasemask exposure**

S. Tonchev<sup>1</sup>, Y. Jourlin<sup>1</sup>, C. Veillas<sup>1</sup>, O. Parriaux<sup>1</sup>, J. Laukkanen<sup>2</sup>, M. Kuittinen<sup>2</sup>; <sup>1</sup>University of Lyon, Lab. H. Curien UMR CNRS 5516 (FR); <sup>2</sup>University of Eastern Finland, Physics and Maths Dept (FI).

A periodic grating with integer number of periods is fabricated at the resist-coated wall of a cylinder by exposing a radial phasemask grating to an axial, radially polarized exposure beam. The created interferogram performs a projection of the planar 2D-space of the high spatial coherence e-beam phasemask onto a cylindrical surface in the 3Dspace. [4855]

NOTES

18:00 - 19:30 Poster session in the foyer

## Senaatszaal, TU Delft

08:30 - 10:00

## SESSION IX

Chair: Kimio Tatsuno, OITDA / Hitachi Ltd. (JP)

NOTES

08:30 - 09:00

Invited talk

**All optical switchable diffraction gratings realized in liquid crystalline composite structure with metamaterial applications**

C. Umeton<sup>1</sup>, L. De Sio<sup>1</sup>, R. Caputo<sup>1</sup>, S. Serak<sup>2</sup>, N. Tabiryani<sup>2</sup>; <sup>1</sup>LICRYL (Liquid Crystals Laboratory, IPCF-CNR) Center of Excellence CEMIF.CAL, Department of Physics, University of Calabria (IT); <sup>2</sup>Beam Engineering for Advanced Measurements Company Winter Park (US).

We report on the fabrication and all optical characterization of two different kinds of switchable diffraction gratings containing light responsive Liquid Crystals. We foresee that, when doped with gold nanoparticles, these structures can be exploited to obtain the tuning of "active" plasmonic responses, to be used for metamaterial applications. [4901]

09:00 - 09:20

**Polarization-selective extraordinary transmission in inductive and capacitive goldnanogrids**

B. Bai<sup>1,2</sup>, X. Li<sup>1</sup>, I. Vartiainen<sup>2</sup>, J. Laukkanen<sup>2</sup>, A. Lehmuskero<sup>2</sup>, J. Turunen<sup>2</sup>; <sup>1</sup>State Key Laboratory of Precision Measurement Technology and Instruments, Department of Precision Instruments, Tsinghua University (CN); <sup>2</sup>University of Eastern Finland, Department of Physics and Mathematics (FI).

We report both theoretically and experimentally that properly designed inductive and capacitive gold nanogrids can exhibit a polarization-selective enhanced-suppressed transmission effect. Exceptionally, in a sparse array of gold nanoparticle chains, an anomalous polarization-dependent resonant suppressed transmission is discovered. The physical mechanisms of these effects are revealed and thoroughly studied and potential applications, such as novel-type compact polarizing filters, ultrasensitive biosensors, and enhanced nonlinear nanostructured devices, are suggested. [4889]

09:20 - 09:40

STUDENT PRESENTATION

**Fabrication of Terahertz Wire-Grid Polarizers**

A. Partanen<sup>1</sup>, J. Väyrynen<sup>2</sup>, S. Hassinen<sup>2</sup>, H. Tuovinen<sup>1</sup>, J. Mutanen<sup>1</sup>, T. Itkonen<sup>1</sup>, P. Silfsten<sup>1</sup>, B. P. Pääkkönen<sup>1</sup>, M. Kuittinen<sup>1</sup>, K. Mönkkönen<sup>2</sup>, T. Venäläinen<sup>3</sup>; <sup>1</sup>University of Eastern Finland, Department of Physics and Mathematics (FI); <sup>2</sup>North Karelia University of Applied Sciences (FI); <sup>3</sup>University of Eastern Finland, Department of Chemistry (FI).

This study shows fabrication of wire-grid grating polarizers for THz region in two ways: by ultra precision diamond machining and by silicon wet etching. The pattern is then transferred to COC and TPX by hot embossing. Aluminum is evaporated on top of the plastic to form the grid. Polarizers were tested with FTIR-spectrophotometer. [4849]

09:40 - 10:00

**TE/TM phase matching for the shallowest half-wave 0th order transmission grating**

T. Kaempfe<sup>1</sup>, S. Tonchev<sup>1</sup>, R. Torrez<sup>2</sup>, J. Lopez<sup>2</sup>, O. Parriaux<sup>1</sup>; <sup>1</sup>Univ. of Lyon, Lab. H. Curien UMR CNRS 5516 (FR); <sup>2</sup>ALPhANOV (FR).

The TE and TM round trip phases in a high index contrast subwavelength binary grating are set to the smallest possible integer numbers of  $2\pi$  and their straight-through phase difference to  $\pi$  by tailoring the reflection and transmission phases at the boundaries of a relatively large period grating for obtaining a half-wave element of close to unity transmission. This TE/TM phase matching principle is applied to linear to radial/azimuthal polarization transformation in the 1030-1064 nm wavelength range with solar cell amorphous silicon as the grating layer material. [4859]

10:00 - 10:30 Coffee break

10:30 - 12:10

## SESSION X

Chair: Sylvania Pereira, Delft University of Technology (NL)

10:30 - 10:50

STUDENT PRESENTATION

**Surface-relief gratings for solar and lighting applications**

T.M. de Jong<sup>1</sup>, D.K.G. de Boer<sup>2</sup>, C.W.M. Bastiaansen<sup>1,3</sup>; <sup>1</sup>Eindhoven University of Technology, Functional Organic Materials & Devices (NL); <sup>2</sup>Philips Research Europe (NL); <sup>3</sup>Queen Mary University of London, Department of Materials (GB).

Surface-relief gratings with sub-wavelength periods can be used to diffract light into or out of a light guide. We show that efficient in- and out-coupling occurs for specific regions in the space of incident angles and discuss how this can be used to design compact solar concentrators and lighting systems. [4908]

Senaatszaal, TU Delft

10:50 - 11:10

STUDENT PRESENTATION

**Metallic nanostructures prepared by nanoimprint lithography for application in solar cells**

*S. Jüchter<sup>1</sup>, H. Hauser<sup>1</sup>, Ch. Wellens<sup>1</sup>, M. Peters<sup>1</sup>, J.C. Goldschmidt<sup>1</sup>, U.T. Schwarz<sup>2</sup>, B. Bläsi<sup>1</sup>;*

*<sup>1</sup>Fraunhofer Institute for Solar Energy Systems ISE (DE); <sup>2</sup>Fraunhofer Institute for Applied Solid State Physics IAF (DE).*

In this study we present first results concerning the advantage of using plasmonic effects in metallic nanostructures for improved photon management in solar cells. [4864]

11:10 - 11:30

**Light scattering of superparamagnetic nanoparticles for biosensing applications**

*J. Schleipen<sup>1</sup>, A. Ranzoni<sup>1</sup>, J. van Kemenade<sup>1</sup>, M. Prins<sup>1</sup>, V. Schmidt<sup>2</sup>, T. Wriedt<sup>2</sup>;* *<sup>1</sup>Philips Research Eindhoven (NL); <sup>2</sup>University of Bremen (DE).*

Biomedical research delivers detailed knowledge about how specific molecules relate to human health and disease. Such specific molecules are called biomarkers when they can be used for diagnostic purposes. For example, the presence of certain protein biomarkers in blood indicates the onset of a heart attack. To enable wide-spread application of biomarker testing, biosensor systems are needed which can easily be used by doctors and patients. The requirements for these point-of-care biosensors are challenging: the systems need to be highly sensitive, reliable, cost-effective, rapid, and very easy to use. [4977]

11:30 - 11:50

STUDENT PRESENTATION

**Enhancing the accuracy of a Hartmann Wavefront Sensor by retrieving the sub-aperture phase distribution**

*A. Polo, F. Bociort, S.F. Pereira, H.P. Urbach;* *Delft University of Technology, Department of Imaging Science and Technology, Optics Research Group (NL).*

A phase retrieval algorithm is applied to the intensity pattern of a Hartmann wavefront sensor to retrieve the phase distribution inside the sub-aperture of a Hartmann hole array. This procedure permits to obtain a rms wavefront reconstruction error one order of magnitude smaller than the one obtained with a typical centroid algorithm. [4884]

11:50 - 12:10

**Novel Fabrication Method of Computational Rainbow Holograms**

*S. Mader<sup>1,2</sup>, T. Kozacki<sup>1</sup>, H. Walter<sup>2</sup>;* *<sup>1</sup>Warsaw University of Technology (PL); <sup>2</sup>OVD Kinegram AG, Optical Technology (CH).*

White-light reconstruction of conventional computational rainbow holograms is limited by the hologram's spatial resolution. A novel method is proposed, theoretically analyzed, and experimentally demonstrated to overcome the reconstruction limitations of computational rainbow holograms of objects at intermediate distances. [4927]

12:10 - 14:00 Lunch break

14:00 - 15:50

**SESSION XI**

*Chair: Cesare Umeton, Università della Calabria (IT)*

14:00 - 14:30

Invited talk

**Theory, experiment, and applications of photonic resonance effects in nanopatterned films-A review**

*R. Magnusson<sup>1,2</sup>, K.J. Lee<sup>1</sup>, J. Yoon<sup>1</sup>, D. Wawro<sup>2</sup>, S. Zimmerman<sup>2</sup>, W. Wu<sup>1</sup>, M. Shokooh-Saremi<sup>1,3</sup>, H.G. Svavarsson<sup>4</sup>, S.H. Song<sup>5</sup>;* *<sup>1</sup>Department of Electrical Engineering, University of Texas at Arlington (US); <sup>2</sup>Resonant Sensors Incorporated (US); <sup>3</sup>Department of Electrical Engineering, Ferdowsi University of Mashhad (IR); <sup>4</sup>School of Science and Engineering, Reykjavik University (IS); <sup>5</sup>Department of Physics, Hanyang University (KR).*

Fundamental photonic resonance effects in thin periodic films with nanoscale features are presented and examples of associated spectral expressions given. Potential applications including biosensors and optical components are discussed. [4972]

14:30 - 14:50

**Avoiding Resonance in Dielectric Reflection Gratings**

*F. Fuchs<sup>1</sup>, D. Michaelis<sup>1</sup>, U.D. Zeitner<sup>1,2</sup>, E.B. Kley<sup>2</sup>;* *<sup>1</sup>Fraunhofer-Institut für Angewandte Optik und Feinmechanik (DE); <sup>2</sup>Institute of Applied Physics, Friedrich-Schiller-University Jena (DE).*

We study the formation of guided mode resonances in multilayer dielectric gratings computationally and devise a scheme of how to avoid them in grating design. [4877]

NOTES



## Senaatszaal, TU Delft

14:50 - 15:10

**Modal representation and normalized scaling rules for ultra narrow-band reflection from a 1D binary corrugation***T. Kämpfe, A.V. Tishchenko, O. Parriaux; University of Lyon, Lab. H. Curien UMR CNRS 5516 (FR).*

A 1D, high index contrast subwavelength binary corrugation on a low index substrate may under normal incidence exhibit arbitrarily narrow band reflection. The conditions for its occurrence are given on the basis of the resonances and interference of two involved grating modes and a phenomenological representation of this effect will be proposed. Normalized heuristic analytical expressions are derived which identify exhaustively all possible structures exhibiting this effect. [4861]

15:10 - 15:30

STUDENT PRESENTATION

**Simultaneous dual wavelength full field imaging in low coherence digital holographic microscopy***Z. Monemhaghdoost<sup>1</sup>, F. Montfort<sup>2</sup>, C. Moser<sup>1</sup>; <sup>1</sup>EPFL, LAPD (CH); <sup>2</sup>Lycée Tec SA (CH).*

We demonstrate a volume diffractive optical element (VDOE) that enables full field of view imaging at two wavelengths in off-axis digital holographic microscopy. Single shot image acquisition at both wavelengths allows for high speed imaging and an increase of the axial unambiguous range from  $0.4\mu\text{m}$  to  $2.5\mu\text{m}$ . [4853]

15:30 - 15:50

STUDENT PRESENTATION

**Effective-medium enhanced three-level grating in resonance domain***M. Oliva<sup>1</sup>, D. Michaelis<sup>1</sup>, F. Fuchs<sup>1</sup>, U.D. Zeitner<sup>1,2</sup>; <sup>1</sup>Fraunhofer Institute for Applied Optics & Precision Engineering (DE); <sup>2</sup>Institute for Applied Physics, Friedrich-Schiller-University Jena (DE).*

A broadband blazed grating with maximum efficiencies above 90% composed of an effective medium structure and a subsequent mode conversion layer is designed for normal incidence in the deep resonance domain. Due to its demanding geometry, a standard multilevel fabrication process requires further optimizations. [4886]

15:50 - 16:30 Coffee break &amp; Whiskey tasting

**17:00 - 18:30 Social programme****Meeting time:** 16:45**Meeting point:** Conference Centre**Venue:** Royal Delft, Koninklijke Porceleyne Fles  
Rotterdamseweg 196**18:30 - 22:30 Conference dinner****Venue:** Partycentrum - Restaurant de Brasserie  
Brasserskade 2a

NOTES

NOTES

Senaatszaal, TU Delft

08:30 - 10:00

SESSION XII

Chair: Joseph Braat, Delft University of Technology (NL)

08:30 - 09:00

Invited talk

**Polarization control with artificial chiral structures**

*M. Kuwata-Gonokami; Department of Physics, Graduate School of Science, Photon Science Center, University of Tokyo (JP).*

Artificial sub-wavelength two dimensional grating structures without mirror symmetry exhibit strong optical activity and rotate polarization states of propagating optical and terahertz waves. We also demonstrate circularly polarized spontaneous emission of quantum dots embedded in a semiconductor chiral photonic crystal. [4867]

09:00 - 09:20

**Calculation of local absorption in three-dimensional structures using the RCWA**

*M. Auer, K.-H. Brenner; University of Heidelberg (DE).*

The calculation of local absorption has become very relevant in the optimization of photodetectors and solar cells. While total absorption is readily available in existing tools, local absorption requires additional effort. Especially in the RCWA, the field calculation is not unique and has significant effects on local absorption. [4924]

09:20 - 09:40

**Optimization of local absorption in layered media**

*K.-H. Brenner; University of Heidelberg (DE).*

For the design of detectors or solar cells, not only the total amount of absorbed power is relevant, but also whether the strongest absorption occurs in the desired layer or in the substrate. An exact definition for local absorption recently has been given. It enables a closed description and thus an optimization of performance. [4943]

09:40 - 10:00

STUDENT PRESENTATION

**Deformable micromirror array for dynamic laser beam shaping and homogenizing**

*J. Masson<sup>1</sup>, R. Bitterl<sup>1</sup>, A. Bich<sup>2</sup>, W. Noell<sup>1</sup>, R. Voelke<sup>2</sup>, K. Weible<sup>2</sup>, N.F. de Rooij<sup>1</sup>; <sup>1</sup>EPFL, SAMLAB (CH); <sup>2</sup>SUSS MicroOptics SA (CH).*

We present a dynamic laser beam shaper that can generate smooth flat-top and Gaussian intensity profiles. It consists of a 100% fill factor membrane, supported by beams or posts, which deforms dynamically to shape and smooth coherent light. The mirror array is fabricated over a scanning stage that enables interference averaging. [4882]

10:00 - 10:30 Coffee break

10:30 - 12:20

SESSION XIII

Chair: Oliver Parriaux, University of Lyon (FR)

10:30 - 11:00

Invited talk

**Modified quantum dot emission with arrays of plasmonic nanoparticles**

*G. Lozano<sup>1</sup>, S.R.K. Rodriguez<sup>1</sup>, M. Verschuuren<sup>2</sup>, R. Gomes<sup>3</sup>, Z. Hens<sup>3</sup>, J. Gómez Rivas<sup>1,4</sup>; <sup>1</sup>Center for Nanophotonics, FOM Institute AMOLF (NL); <sup>2</sup>Philips Research Laboratories (NL); <sup>3</sup>Physics and Chemistry of Nanostructures, Center for Nano and Biophotonics, Ghent University (BE); <sup>4</sup>COBRA Research Institute, Eindhoven University of Technology (NL).*

We present an experimental study of the modified emission of a layer of quantum dots (QDs), just a few nanometers thick, with plasmonic structures. The coupling of localized surface plasmon resonances in metallic particles through in-plane diffraction in arrays of these particles leads to the formation of quasi-bound surface states known as surface lattice resonances (SLRs). The recent observation of extremely narrow SLRs in arrays of gold nanoparticles has created expectations for these arrays as plasmonics sensors. We show here that these resonances can lead to a strong modification of the emission due to the modified density of optical states at which the excited QDs can decay and the diffractive coupling of SLRs into the far-field. [4957]

NOTES

## Senaatszaal, TU Delft

11:00 - 11:20

STUDENT PRESENTATION

NOTES

**In-situ hologram correction in SLM setup**

*V. Carrat, B. Viaris de Lesegno, L. Pruvost; Laboratoire Aimé Cotton CNRS UPR3321 Bat 505 Univ. Paris-Sud (FR).*

By adding another wavelength in an existing SLM phase modulation setup, we developed a robust method to get the birefringence map of the modulator. By this way we are able to see and correct the defects in-situ without disturbing the running experiment. [4902]

11:20 - 11:40

**Hybrid diffractive-refractive wide field of view objective**

*E. Sokolova<sup>1</sup>, I. Livshits<sup>2</sup>; <sup>1</sup>River Diagnostics B.V. (NL); <sup>2</sup>Lab CAD of opto-information and energy saving systems, NRU ITMO (RU).*

The method of substitution lenses with complicated high order aspherical surfaces by the hybrid diffractive-refractive lenses is proposed. Two examples with the same number of elements are given. As a result the image quality is improved without using aspheric surfaces. [4866]

11:40 - 12:00

**Unidirectional Surface Plasmon Launchers and Decouplers**

*A. Baron<sup>1</sup>, E. Devaux<sup>2</sup>, J.-C. Rodier<sup>1</sup>, J.-P. Hugonin<sup>1</sup>, E. Rousseau<sup>1</sup>, C. Genet<sup>2</sup>, T.W. Ebbesen<sup>2</sup>, P. Lalanne<sup>1,3</sup>; <sup>1</sup>Laboratoire Charles Fabry, Institut d'Optique, Univ Paris-Sud, CNRS (FR); <sup>2</sup>ISIS, Université de Strasbourg et CNRS (UMR 7006) (FR); <sup>3</sup>Lab. de Photonique Numérique et Nanosciences, Institut d'Optique, Univ Bordeaux 1, CNRS (FR).*

We designed, fabricated and characterized a compact novel device composed of eleven grooves which displays unprecedented performances regarding two important figures-of-merit of SPP launchers, namely the launching/decoupling efficiencies and the extinction ratio that characterizes the directionality of the launching. [4851]

12:00 - 12:20

**Compressive sensing inverse imaging techniques in diffraction optics**

*V. Katkovnik, A. Migukin, J. Astola; Department of Signal Processing, Tampere University of Technology (FI).*

Let us start from a wave field reconstruction in the following simple setup. A coherent complex-valued wave field propagates from an object plane to a parallel image plane. This free space forward propagation is modeled by the Rayleigh-Sommerfeld diffraction integral. The problem is to reconstruct the complexvalued wave field at the object plane from complex-valued observations given at the image plane. [4824]

12:20 END OF EOS TOPICAL MEETING

NOTES

Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

4764\_DO2012\_01

STUDENT PRESENTATION

**Plasmonic nanosensor in the treatment of cancer: An attempt to conquer the immortal illness***S. Das, J. Turunen; University of Eastern Finland, Department of Physics and Mathematics (FI).*

We present a novel method based on silver nanoparticle-generated transient photothmal vapour nanobubbles which is effective in the diagnosis (by optical scattering) and treatment (by mechanical, nonthermal and selective destruction of target cells) of cancerous cells.

4828\_DO2012\_03

STUDENT PRESENTATION

**Accurate measurement of the Bidirectional Transmittance Distribution Function of Holographic Diffusers***J. Audenaert<sup>1</sup>, G. Durinck<sup>1</sup>, F. Leloup<sup>1</sup>, G. Deconinck<sup>2</sup>, P. Hanselaer<sup>1,2</sup>; <sup>1</sup>Light&Lighting Laboratory-Catholic University College Sint-Lieven (BE); <sup>2</sup>ESAT-ELECTA – Katholieke Universiteit Leuven (BE).*

Due to the instrument characteristics of the measurement device, large deviations can be encountered between theoretical and experimental Bidirectional Transmittance Distribution Functions. The application of an iterative algorithm to simulated and measured BTDFs, based on the theorem of Bayes, is proposed. A significant improvement of the accuracy of the distribution functions is achieved compared to the raw untreated data.

4832\_DO2012\_04

**Hybrid diffractive-refractive objective for Raman microscopy/spectroscopy***E. Sokolova; River Diagnostics B.V. (NL).*

A number of variants of a compact microscope objective all lenses of which are made of the same material (preferably fused silica) are presented. One of the lens surfaces (plane or spherical) is diffractive. Other lens surfaces are spherical.

4836\_DO2012\_05

STUDENT PRESENTATION

**Definition of a compact infrared camera based on a binary implemented continuously self-imaging grating***M. Piponnier<sup>1</sup>, G. Druart<sup>1</sup>, N. Guérineau<sup>1</sup>, J.-L. De Bougrene<sup>2</sup>, J. Primot<sup>1</sup>; <sup>1</sup>ONERA – The French Aerospace Lab (FR); <sup>2</sup>Telecom Bretagne, Optics Department (FR).*

A dewar is a sealed environment used to cool infrared detectors. Specific diffractive gratings such as axicons or continuously self-imaging gratings are investigated to integrate a simple imagery function inside the dewar, because they are known to have a long depth of focus and a small mass.

4837\_DO2012\_06

**Diffractive and interferometric methods to characterize photopolymers with crystal liquid molecules as holographic recording material***S. Gallego, A. Márquez, M. Ortuño, J. Francés, P. Gallego, A. Beléndez, I. Pascual; Universidad de Alicante, Instituto Universitario de Física Aplicada a las Ciencias y las Tecnologías (ES).*

We present two methods, interferometry at the zero spatial frequency limit and analysis of diffracted orders for very low spatial frequency gratings, to characterize photopolymers with dispersed nematic liquid crystals. These methods provide us information in real time about the transformations taking place inside the material during recording.

4848\_DO2012\_07

**Theoretical description of a 2D volume hologram with sources***Y.G. Boucher<sup>1,2</sup>, F.F.L. Bentivegna<sup>3</sup>, D. Moussa Djama<sup>3</sup>, T.V. Galstian<sup>4</sup>; <sup>1</sup>Université Européenne de Bretagne (UEB) (FR); <sup>2</sup>CNRS, UMR 6082 Foton (FR); <sup>3</sup>RESO, École Nationale d'Ingénieurs de Brest (FR); <sup>4</sup>COPL, Laval University (CA).*

We present an all-analytical description of a two-dimensional active volume hologram consisting of a distributed cavity with localized emitters. Using both coupled-mode theory and extended matrix formalism, we show that the modal response of the structure is fully determined by a set of dimensionless parameters.

4850\_DO2012\_08

**Application of Fourier modal method to structures including negative-index media***H. Ichikawa; Ehime University, Faculty of Engineering (JP).*

Fourier modal method is applied to analysis of optical elements which include negative-index media. Its implementation and example computation are presented. It is shown that the method can handle negative-index media, but is associated with some problems.

4852\_DO2012\_09

STUDENT PRESENTATION

**Double DOE system for near-field imaging formation***J.M. Herrera-Fernandez, L.M. Sanchez-Brea, E. Bernabeu; Universidad Complutense de Madrid, Optics Department, Applied Optics Complutense Group (ES).*

We present an algorithm for forming the desired image in the near field with two DOEs; the first DOE modulates the amplitude and it is equal to the target image, whereas the second DOE modulates the phase.

NOTES

Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

4854\_DO2012\_10

**Stokes parameters for diffraction vector fields**

*J. Silva-Barranco<sup>1</sup>, P. Martínez Vara<sup>2</sup>, T. Quarton<sup>3</sup>, G. Martínez-Niconoff<sup>1</sup>; <sup>1</sup>Optics Department, Instituto Nacional de Astrofísica, Óptica y Electrónica (MX); <sup>2</sup>Benemerita Universidad Autónoma de Puebla, BUAP, Dpto de Ingeniería (MX); <sup>3</sup>University of Texas at Dallas UTD, Physics Department (US).*

We define the vector correlation function between two amplitude functions mutually orthogonal. We show that this function satisfies the Helmholtz equation and it allows the Stokes parameters for arbitrary diffraction fields to be obtained.

4856\_DO2012\_11

**Exact modelling of large pixelated DOEs with wavelength-scale feature size**

*T. Kaempfe, A. Tishchenko; University of Lyon, Lab. H. Curien UMR CNRS 5516 (FR).*

An ultra-fast exact electromagnetic modeling approach based on the generalized source method (GSM) achieving  $O(N \log N)$  calculation time and memory resort is applied to the simulation of the diffraction of a relatively large section of a 2D pixelated DOE in the form of square pillars of identical cross-section. The size of the considered area is several wavelengths across, enough to achieve a reasonable separation of the central region from the influence of neighbouring areas, enabling the use of a stitching method for the simulation of very large elements that could so far only be modeled by scalar approximations. A time and memory resort comparison is made between a standard scalar method and the rigorous GSM-based simulation of the spectra of the element.

4857\_DO2012\_12

**Fabrication of (sub)wavelength resonant DOEs: a come-back of wet etching**

*S. Tonchev, O. Parriaux; University of Lyon, Lab. H. Curien UMR CNRS 5516 (FR).*

Resonant diffractive elements are characterized by a strong field enhancement in the neighbourhood of the surface corrugation due to the presence of a surface wave. This confers a large radiation strength to a shallow corrugation which can therefore be wet-etched with negligible underetch. Low cost, high manufacturing productivity is expectable.

4863\_DO2012\_13

**Accuracy analysis of lamellar diffraction gratings by means of simplified theories**

*J. Francés, S. Gallego, S. Bleda, C. Neipp, A. Márquez, I. Pascual, A. Beléndez; Universidad de Alicante, Instituto Universitario de Física Aplicada a las Ciencias y las Tecnologías (ES).*

The validity of both the scalar diffraction theory and the effective medium theory is evaluated by the comparison of the diffraction efficiencies predicted from both simplified theories to accurate results calculated by the finite-difference time-domain method.

4868\_DO2012\_15

STUDENT PRESENTATION

**Analysis of diffractive optical elements generated onto photopolymer materials using a liquid crystal display**

*A. Márquez<sup>1,3</sup>, S. Gallego<sup>1,3</sup>, M. Riquelme<sup>1</sup>, M. Ortuño<sup>1,3</sup>, M.L. Álvarez<sup>1,3</sup>, A. Beléndez<sup>1,3</sup>, I. Pascual<sup>2,3</sup>; <sup>1</sup>Dpt. de Física, Ing. de Sistemas y Teoría de la Señal, Univ. de Alicante (ES); <sup>2</sup>Dpt. de Óptica, Farmacología y Anatomía, Univ. de Alicante (ES); <sup>3</sup>I.U. Física Aplicada a las Ciencias y las Tecnologías, Univ. de Alicante (ES).*

We produce phase-only diffractive optical elements (DOE) onto photopolymer. To this goal we use a liquid crystal display (LCD), displaying the DOE amplitude master, which is imaged onto the photopolymer. Various challenges need to be overcome such as accurate combined control of LCD and photopolymer and proper imaging of the master onto the photopolymer plane, which will be presented in the paper.

NOTES

Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

4870\_DO2012\_16

**Near field of cylindrical gratings with point source illumination***F.J. Torcal-Milla, F.J. Salgado-Remacha, L.M. Sanchez-Brea, E. Bernabeu; Universidad Complutense de Madrid, Optics Department (ES).*

We analyze how a diffraction grating engraved on the external surface of a cylinder diffracts the light in the near field. We assume a point source illuminating the grating and show how the distance between the source and the grating affects to the diffracted intensity. There is a situation in which the curvature is compensated giving parallel fringes.

4872\_DO2012\_17

**General properties of metallic gratings with subwavelength indentations***P. Lalanne<sup>1,2</sup>, H.T. Liu<sup>3</sup>; <sup>1</sup>Laboratoire Charles Fabry, Institut d'Optique, Univ Paris-Sud, CNRS (FR); <sup>2</sup>Lab. de Photonique Numérique et Nanosciences, Institut d'Optique (FR); <sup>3</sup>Key Laboratory of Optical Information Science and Technology, Ministry of Education, Institute of Modern Optics, Nankai University (CN).*

Under illumination by a transverse-magnetic (TM) wave, the electromagnetic field scattered on metallic surfaces corrugated by one-dimensional periodic arrays of tiny indentations exhibits several general properties. In general, the field on the surface between the indentations does not contain any surface-plasmon-polariton (SPP). Exception occurs in a narrow energy band centered around  $\lambda = \lambda_{sp}$ , the SPP-phased matched wavelength obtained when the momentum of the incident wave matches the SPP momentum of the *flat* interface.

4873\_DO2012\_18

STUDENT PRESENTATION

**Generalized Phase Contrast with matched filtering using LCoS pico-projectors***A. Bañas, D. Palima, J. Glückstad; DTU Fotonik, Department of Photonics Engineering Programmable Phase Optics Technical University of Denmark (DK).*

We report a beam shaping system for generating high intensity programmable optical spots using mGPC: matched filtering combined with Generalized Phase Contrast applying two consumer handheld pico-projectors. Such a system presents a low cost alternative for optical trapping and manipulation, optical lattices and other beam shaping applications usually implemented with high-end spatial light modulators.

4876\_DO2012\_19

**Femtosecond laser pulse ablation with two-dimensional two-grating interferometer***J.J.J. Kaakkunen, M. Silvennoinen, K. Päiväsaari, P. Vahimaa; University of Eastern Finland, Department of Physics and Mathematics (FI).*

In this paper, the realization of four waves interference pattern using two-dimensional two-grating interferometer is presented. Interference pattern is used to ablate large areas of under micron feature size structures into various materials using Ti:Sapphire femtosecond laser with a central wavelength of 800 nm.

4897\_DO2012\_20

**Experimental and theoretical investigations of nanosecond laser damage on diffractive optics***R. Burla<sup>1</sup>, X. Fu<sup>1</sup>, F. Wagner<sup>1</sup>, G. Demesy<sup>1</sup>, L. Gallais<sup>1</sup>, J.Y. Natoli<sup>1</sup>, M. Commandré<sup>1</sup>, S. Gautier<sup>2</sup>, S. Tisserand<sup>2</sup>; <sup>1</sup>Institut Fresnel, CNRS, Aix-Marseille Université (FR); <sup>2</sup>SILIOS TECHNOLOGIES, Z.I. Peynier-Rousset (FR).*

The laser damage resistance of phase mirrors used for laser beam shaping in high peak power laser applications is studied. Laser Induced Damage Thresholds measurements are done at 1064nm, 6ns and the results are compared with simulations of the electric field within the multi-layer by using a wave propagation computer code.

4888\_DO2012\_22

**Wideband, wide angular spectrum resonant reflection by mode coalescence in dual-mode slab waveguide***Y. Jourlin<sup>1</sup>, S. Tonchev<sup>1</sup>, A.V. Tishchenko<sup>1</sup>, C. Pédrix<sup>1</sup>, O. Parriaux<sup>1</sup>, D. Jamon<sup>2</sup>, F. Lacour<sup>3</sup>; <sup>1</sup>University of Lyon, Lab. H. Curien UMR CNRS 5516 (FR); <sup>2</sup>University of Lyon, Lab. Telecom. C. Chappe (FR); <sup>3</sup>Eurofarad (FR).*

A deep grating in a dual-mode high index waveguide causes the coalescence of the two reflection peaks into a single peak of broader wavelength spectrum and high angular robustness. This property extends the applicability of resonant reflection to the wide field of light processing systems dealing with low spatial and temporal coherence beams. Hydrogenated amorphous silicon deposited on polymer at 1500C is used as the low-loss slab-waveguide in the near-IR wavelength range.

NOTES

Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

4892\_DO2012\_23

STUDENT PRESENTATION

**Fabrication of Linear Grating like Laser Induced Periodic Surface Structures**

*M.V.J. Silvennoinen, J.J.J. Kaakkunen, K. Paivasaari, P. Vahimaa; University of Eastern Finland, Department of Physics and Mathematics (FI).*

Fabrication of the Laser Induced Periodic Surface Structures using IR femtosecond laser pulses is studied. It has been noticed that by using LCOS-spatial light modulator it is possible to fabricate large areas of linear grating like structures. The spatial light modulator can be used also to focus the pulses using a hologram of a cylindrical lens.

NOTES

4894\_DO2012\_24

STUDENT PRESENTATION

**Influence of phase fluctuations of a LCOS based Hartmann-Shack sensor**

*J.L. Martínez-Fuentes<sup>1</sup>, J. Campos<sup>2</sup>, I. Moreno<sup>1</sup>, A. Vargas<sup>3</sup>; <sup>1</sup>Universidad Miguel Hernández de Elche, Dept. de Ciencia de Materiales, Óptica y Tecnología Electrónica (ES); <sup>2</sup>Universitat Autònoma de Barcelona, Departament de Física (ES); <sup>3</sup>Universidad de la Frontera, Departamento de Ciencias Físicas (CL).*

We study a Hartmann-Shack sensor implemented by means of a parallel aligned LCOS microdisplay presenting temporal phase fluctuations and a CCD camera. Concretely, we perform a comparative analysis of the sensing position stability versus the camera integrating time and the size of the lenses addressed to the display.

4895\_DO2012\_25

STUDENT PRESENTATION

**Controlling plasmon resonance peaks in gold-based metamaterials**

*J. Lehtolahti<sup>1</sup>, J. Laukkanen<sup>1</sup>, M. Kauranen<sup>2</sup>, M. Kuittinen<sup>1</sup>; <sup>1</sup>University of Eastern Finland, Department of Physics and Mathematics (FI); <sup>2</sup>Tampere University of Technology, Department of Physics (FI).*

By varying the sizes and distances of circular subwavelength gold nanoparticles in two-dimensional periodic arrays it is possible to adjust both the plasmon resonance peak wavelengths and the strength of the resonances, as our numerical calculations show.

4896\_DO2012\_26

STUDENT PRESENTATION

**Twin-SLM Complex Light Modulation for Computer Holography**

*A. Czerwinski<sup>1</sup>, M. Makowski<sup>1</sup>, J. Suszek<sup>1</sup>, A. Siemion<sup>1</sup>, M. Sypek<sup>1</sup>, Z. Jaroszewicz<sup>2</sup>, A. Kolodziejczyk<sup>1</sup>; <sup>1</sup>Warsaw University of Technology, Faculty of Physics (PL); <sup>2</sup>Institute of Applied Optics (PL).*

We present an experimental evaluation of the technique of a complex modulation of the amplitude and phase of laser light. We use a pair of identical spatial Light Modulators (SLM) manufactured in the LCoS technology (Liquid Crystal on Silicon). The evaluation is performed by addressing the SLMs with computer-generated holograms, which is aimed at future applications in a compact, lensless projection of color images.

4898\_DO2012\_27

**Partially coherent beams with spatially varying correlations**

*H. Lajunen, T. Saastamoinen; University of Eastern Finland, Department of Physics and Mathematics (FI).*

We study a new class of partially coherent beams with spatially varying correlations. It is shown by numerical simulations that such inhomogeneous coherence distributions may produce extraordinary propagation-induced changes, such as sharpened and laterally shifted maxima of the beam intensity.

4899\_DO2012\_28

**Liquid-crystal-based diffractive optical elements**

*S. Valyukh<sup>1</sup>, P. Tytarenko<sup>2</sup>; <sup>1</sup>Laboratory of Applied Optics, Department of Physics, Chemistry and Biology (IFM), Linköping University (SE); <sup>2</sup>Institute of Semiconductor Physics (UA).*

Tunable liquid-crystal (LC) -based diffractive optical elements (DOE) are studied. We investigate limitations and potential functional abilities of LC DOE utilizing homogeneous and inhomogeneous alignment. Influence of the anchoring energy of LC on diffractive properties of DOE is discussed.

4905\_DO2012\_29

STUDENT PRESENTATION

**Detection of dust and scratches on photographic material by combining dark field illumination and crossed polarization**

*G. Trumpp, A. Wassmer, R. Gschwind; University of Basel, Imaging & Media Lab (CH).*

A solution for the elimination of dust and scratches on transparent photographic material is investigated. We are designing a scanning system that will detect defects taking advantage of depolarization of light scattered by irregularities in the film surface. This phenomenon is studied analytically to define the optimal optical setup.

Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

NOTES

4906\_DO2012\_30

**Diffraction pattern from mature and immature red blood cells**

*M. Mihailescu<sup>1,2</sup>, A. Gheorghiu<sup>2</sup>, J. Costescu<sup>2</sup>; <sup>1</sup>National Institute for Research and Development in Microtechnologies (RO); <sup>2</sup>Politehnica University from Bucharest (RO).*

We present our study regarding mature and immature red blood cells diffraction patterns in Fresnel and Fraunhofer approximation. We model their shapes taking into account dimensions established experimentally in digital off-axis holographic microscopy. We considered few types: mature and immature, deformed and undeformed red blood cells.

4909\_DO2012\_31

STUDENT PRESENTATION

**Beam shaping and light focusing by active selfoc microlenses**

*A.I. Gomez-Varela<sup>1</sup>, M.T. Flores-Arias<sup>1</sup>, C. Bao-Varela<sup>1</sup>, X. de la Fuente<sup>2</sup>, C. Gomez-Reino<sup>1</sup>;*

*<sup>1</sup>Grupo de "Microóptica y Óptica GRIN", Unidad Asociada al Instituto de Ciencias de Materiales de Aragón, ICMA/CSIC, Zaragoza, y Facultade de Óptica e Optometría, Universidade de Santiago (ES); <sup>2</sup>Instituto de Ciencia de Materiales de Aragón (Universidad de Zaragoza-CSIC) (ES).*

Light propagation in active GRIN (GRAdient-INdex) materials with gain or loss is studied. Results are applied to an active selfoc microlens in order to examine Gaussian beam transformation. In particular, beam half-width and curvature radius are analyzed to find beam shaping and light focusing conditions.

4910\_DO2012\_32

STUDENT PRESENTATION

**Geometric metrology of gold nanoparticle ensembles with optical extinction spectroscopy**

*N. Xu, B. Bai, Q. Tan, G. Jin; State Key Laboratory of Precision Measurement Technology and Instruments, Department of Precision Instruments, Tsinghua University (CN).*

We present a method based on optical extinction spectroscopy to fast characterize the geometric features (including size, aspect ratio, and geometric distribution) of gold nanosphere and nanorod ensembles. The results are compared with those obtained by transmission electron microscopy, showing the effectiveness of the method. The influence of the particle parameters is analyzed and the possible methods for solving the ill-posed inverse problem are discussed.

4912\_DO2012\_33

**Wavelength-Dispersive Micro Prism with Periodic Metal Structure in a Silicon Slab-Waveguide**

*S. Kameda, T. Otsuka, T. Morikawa, A. Mizutani, H. Kikuta; Osaka Prefecture University, School of Engineering (JP).*

A micro triangular prism with periodic metal structure was fabricated in a silicon slab waveguide for a wavelength-dispersive device by use of the electroless copper plating. The micro prism was 1.2 $\mu\text{m}$  thick and the structure period was 0.3 $\mu\text{m}$ . The light refraction by the micro prism was observed for the 1.55 $\mu\text{m}$  wavelength light.

4913\_DO2012\_34

STUDENT PRESENTATION

**Fabrication of metallic optical elements embedded in dielectric substrates by the bottom-up fill electroless copper plating**

*T. Otsuka<sup>1</sup>, T. Morikawa<sup>1</sup>, S. Kameda<sup>1</sup>, A. Mizutani<sup>1</sup>, H. Kikuta<sup>1</sup>, S. Shingubara<sup>2</sup>; <sup>1</sup>Osaka Prefecture University, Graduate School of Engineering (JP); <sup>2</sup>Kansai University, Faculty of Engineering Science (JP).*

Optical elements with metal-and-dielectric structure were fabricated by use of the "bottom-up fill" electroless copper plating. The 200nm-wide and 1200nm-deep trenches were filled up with copper. A wire-grid polarizer for near-infrared light and a micro-prism in the silicon-on-insulator substrate were fabricated.

4915\_DO2012\_35

STUDENT PRESENTATION

**Comparison of convolution and Time Dependent Random Phase methods for modelling of the space invariant optical system with spatially incoherent illumination**

*A. Czerwiński, K. Kakarenko, M. Sypek, M. Makowski, I. Ducin, J. Suszek, A. Kołodziejczyk; Warsaw University of Technology, Faculty of Physics (PL).*

In this paper we present the comparison of two methods for modeling space invariant optical system with spatially incoherent quasi-monochromatic illumination. The first one is a convolution method. The second one is a method based on the Time Dependent Random Phase (TDRP) algorithm. Both approaches provide equivalent results.

However, the TDRP method maintains energy relations and magnification of the system. In the convolution method the pre-processing of the input image is required.



Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

4917\_DO2012\_36

**Measurement of aspherical surfaces using CGH test plates**

*A.G. Poleshchuk<sup>1</sup>, R.K. Nasyrov<sup>1</sup>, J.-M. Asfour<sup>2</sup>; <sup>1</sup>Institute of Automation & Electrometry SB RAS (RU); <sup>2</sup>Dioptric GmbH (DE).*

A new method for measuring aspherical surface shapes using computer generated holograms (CGH) as a test plates is presented. Test results of the CGH design, fabrication and measurements are discussed.

4919\_DO2012\_38

STUDENT PRESENTATION

**Fast one-exposure digital holography based on the Talbot effect and phase-shifting method**

*Ag. Siemion, M. Sypek, An. Siemion, J. Suszek, A. Kolodziejczyk, M. Makowski; Warsaw University of Technology, Faculty of Physics (PL).*

Fast one-exposure digital holography technique based on the Talbot effect and phase-shifting method is presented. The optical system is very compact and does not need any imaging elements or advanced cameras. The reconstruction algorithm is simple, quick and based on the optical methods, avoiding any approximation.

4920\_DO2012\_39

**Hybrid Collimating Optics for RGB LEDs**

*T. Bonenberger, J. Baumgart; Hochschule Ravensburg-Weingarten, ZAFH-LED OASYS (DE).*

A technique that has been established to correct color aberrations in photographic optics was used to reduce color fringes in LED lighting applications. This technique is based on the combination of refractive, reflective and diffractive elements realizing a hybrid optical system.

4921\_DO2012\_40

**Focusing properties of the non-paraxial Light Sword Optical Element**

*K. Kakarenko<sup>1</sup>, I. Ducin<sup>1</sup>, Z. Jaroszewicz<sup>2</sup>, M. Makowski<sup>1</sup>, A. Kolodziejczyk<sup>1</sup>, K. Petelczyc<sup>1</sup>, J. Suszek<sup>1</sup>, M. Sypek<sup>1</sup>; <sup>1</sup>Warsaw University of Technology, Faculty of Physics (PL); <sup>2</sup>Institute of Applied Optics and National Institute of Telecommunications (PL).*

We present an analysis of focusing realized by the light sword optical element (LSOE), especially designed for imaging with extended depth of field. The obtained numerical results illustrate that the non-paraxial design of such element can improve its performance in an imaging system.

4923\_DO2012\_41

STUDENT PRESENTATION

**Numerical modeling of the EUV microscope optical setup**

*D. Wojnowski, M. Sypek, J. Suszek, A. Kolodziejczyk; Faculty of Physics, Warsaw University of Technology (PL).*

In this work we present an effective algorithm for modeling the imaging phenomena by the EUV microscope optical setup. Using the algorithm, we focus on source's spectrum influence on spatial resolution of the microscope.

4926\_DO2012\_42

**Diffractive elements for correction of aberrations of white light illumination systems: design and fabrication**

*M. Škereň, P. Fiala, J. Svoboda, J. Hopp, M. Květoň; Czech Technical University in Prague, Faculty of Nuclear Sciences and Physical Engineering, Department of Physical Electronics (CZ).*

In this paper, the application of synthetic diffractive elements is presented for correction of chromatic aberration of white light illumination systems. The cost effective fabrication technology is also presented, which is compatible with the commonly used molding process for fabrication of plastic optical elements.

4929\_DO2012\_43

**Fourier modal analysis of image formation in aplanatic systems**

*H.J. Hyvärinen<sup>1,2</sup>, S. Mehta<sup>3</sup>, J. Turunen<sup>1</sup>; <sup>1</sup>University of Eastern Finland, Department of Physics and Mathematics (FI); <sup>2</sup>Institute of Microelectronics, A\*STAR (SG); <sup>3</sup>Cellular Dynamics Program, Marine Biological Laboratory (US).*

Fourier Modal Method and Abbe theory are applied to analyze coherent image formation of periodic objects in aplanatic imaging systems. The model is applied to bright-field imaging systems, and the results are compared to respective approximative results.

NOTES

4930\_DO2012\_44

**Coherence–polarization mixing in resonance waveguide gratings***T. Saastamoinen, I. Vartiainen, J. Tervo, M. Kuittinen; University of Eastern Finland, Department of Physics and Mathematics (FI).*

We show that resonance gratings can be used to transfer partial spatial correlation to partial polarization even in the case of fully polarized incident beam. This phenomenon is based on the fact that one of the orthogonal polarization components can be coupled into the leaky mode while the other one is reflected without being coupled in the grating.

4931\_DO2012\_45

STUDENT PRESENTATION

**Simulation of broad area laser diodes with elementary mode method***H. Partanen; University of Eastern Finland, Department of Physics and Mathematics (FI).*

Propagation of partially coherent laser beam is modeled with elementary mode method to demonstrate its effectiveness. Intensity and degree of coherence is simulated and compared to results from Wolf's coherent mode representation.

4933\_DO2012\_46

**Fabrication and testing of highly efficient resonance domain diffractive optical elements***O. Barlev<sup>1</sup>, M.A. Golub<sup>1</sup>, A.A. Friesem<sup>2</sup>, D. Mahalu<sup>2</sup>; <sup>1</sup>Tel Aviv University, Dept. of Electrical Engineering (IL); <sup>2</sup>Weizmann Institute of Science, Dept. of Physics of Complex Systems (IL).*

Experimental procedures for recording, evaluating different resonance domain diffractive optical elements are presented. Results reveal that when recorded in fused silica (FS) material as surface relief gratings with a period 520 nm and aspect ratio up to 2.5, they can have very high diffraction efficiency when illuminated with visible light.

4934\_DO2012\_47

STUDENT PRESENTATION

**A short paper on the THz paper optics***A. Siemion<sup>1</sup>, Ag. Siemion<sup>1</sup>, J-L Coutaz<sup>2</sup>, J. Suszek<sup>1</sup>, M. Makowski<sup>1</sup>, I. Ducin<sup>1</sup>, A. Kołodziejczyk<sup>1</sup>, M. Sypek<sup>1</sup>; <sup>1</sup>Warsaw University of Technology, Faculty of Physics (PL); <sup>2</sup>University of Savoie, IMEP-LAHC (FR).*

The diffractive optical element for THz applications made of paper was designed and manufactured. It is a phase only structure, which is transparent for THz radiation. The experimental results by means of Time-Domain Spectroscopy shows bending and focusing of the THz beam (for 0,3-1,5 THz) after the paper element.

4935\_DO2012\_48

**Computer generated matched spatial filters for multiplexing and demultiplexing of spatial modes of coherent light***S. Shwartz, M.A. Golub, S. Ruschin; Tel Aviv University, Department of Electrical Engineering (IL).*

Computer generation of multichannel diffractive optical elements matched to several orthogonal transverse modes of laser beam was solved by a mathematical generating functions concept. Experimental and computer simulated results for reconstruction of modes in coherent optical correlator are in good match.

4936\_DO2012\_49

STUDENT PRESENTATION

**Electromagnetic theory of Hanbury Brown–Twiss effect***T. Hassinen<sup>1</sup>, J. Tervo<sup>1</sup>, T. Setälä<sup>2</sup>, A.T. Friberg<sup>1,2,3</sup>; <sup>1</sup>University of Eastern Finland, Department of Physics and Mathematics (FI); <sup>2</sup>Aalto University, Department of Applied Physics (FI); <sup>3</sup>Royal Institute of Technology (KTH), Department of Microelectronics and Applied Physics (SE).*

We analyze the Hanbury Brown - Twiss experiment with thermal vector waves in the space-frequency domain. It is shown that the spectral degree of electromagnetic coherence fully characterizes the normalized correlation of intensity fluctuations measured in the experiment, a result analogous to the scalar-wave analysis.

4937\_DO2012\_50

STUDENT PRESENTATION

**Simulation and design of photonic crystal cavities for optical trapping of bacteria in water***M.M. van Leest, J.T. Heldens, J. Caro; Delft University of Technology, Kavli Institute of Quantum-Nanoscience (NL).*

A photonic crystal (PhC) cavity can be used as optical trap for single microorganisms and is promising for Raman fingerprinting these objects. We design three PhC cavities for optical trapping experiments, using the 3D FDTD method. Design criteria are a small trapping-induced resonance shift and a high in-plane transmission.

Tuesday, 28 February, 18:00 - 19:30 - Foyer, TU Delft

4941\_DO2012\_51

**Bifocal zone plates designed using the Thue-Morse sequence**

*W.D. Furlan<sup>1</sup>, A. Calatayud<sup>2</sup>, L. Remón<sup>2</sup>, J.A. Monsoriu<sup>2</sup>, P. Andrés<sup>1</sup>; <sup>1</sup>Departamento de Óptica, Universitat de València (ES); <sup>2</sup>Centro de Tecnologías Físicas, Universitat Politècnica de València (ES).*

We present a new family of diffractive lenses constructed using the aperiodic sequence known as the Thue-Morse sequence. The result is a novel kind of bifocal diffracting lenses with self-similar focalization properties. The axial irradiance provided by several members of the family are presented.

4942\_DO2012\_52

**Traveling waves approach for Airy beams**

*H.A. Jiménez-Romero<sup>1</sup>, S. Chávez-Cerda<sup>2</sup>; <sup>1</sup>Depto. De Física Cinvestav-IPN (MX); <sup>2</sup>Instituto Nacional de Astrofísica, Óptica y Electrónica (MX).*

We demonstrate that Airy beams share many physical properties with Bessel beams due to the fact that Airy functions are Bessel functions of order  $\nu=1/3$ . This allows to provide a clear and easy picture for the physics of Airy beams and their properties of propagation. We also show the existence of high order Airy beams.

4946\_DO2012\_53

**Sensitivity analysis of rigorous effects in diffractive null optics**

*S. Peterhänzel, C. Pruss, W. Osten; University Stuttgart, Institut for Applied Optics (DE).*

The increasing demand of precise measurements of steep high-precision free form surfaces leads to higher frequencies in computer generated holograms for the interferometric null test method. In this paper we discuss the impact of phase errors introduced by rigorous effects in diffractive null optics.

4952\_DO2012\_54

**Joint Research on scatterometric characterization of planar and curved diffractive optics**

*A. Lassila<sup>1</sup>, B. Bodermann<sup>2</sup>, J. Turunen<sup>3</sup>, P.-E. Hansen<sup>4</sup>, M. Wurm<sup>2</sup>, S. Siitonen<sup>5</sup>, T. Saastamoinen<sup>3</sup>, V. Korpelainen<sup>1</sup>; <sup>1</sup>Mittatekniikan Keskus (FI); <sup>2</sup>Physikalisch-Technische Bundesanstalt (DE); <sup>3</sup>University of Eastern Finland (FI); <sup>4</sup>Dansk Fundamental Metrolog (DK); <sup>5</sup>Nanocomp Oy Ltd (FI).*

Within a European joint research project it is aimed to develop fast and simple methods for process development and quality control of both diffractive and diffractive-refractive optics in industrial environments. We describe the status of scatterometry for these applications and the intended goals of this project.

4953\_DO2012\_55

**Using the optical code for simulation of the fiber-optic ESPI set-up design**

*V. Abaskin, E. Achimova; Academy of Sciences of Moldova, Institute of Applied Physics, Center of Optoelectronics (MD).*

This article has discussed design of optical part of Electronic Speckle Pattern (ESPI) Interferometer set-up in ZEMAX code. Modulation Transfer Function (MTF), Interferometric Fringe Analysis, Image Analysis, Wavefront Map (WM) was calculated for verifying and predicting the performance of optical part of fiber-optic ESPI set-up.

5008\_DO2012\_56

**Apodized phase masks obtained by means of HEBS glasses**

*T. Osuch<sup>1,2</sup>, A. Kowalik<sup>3</sup>, Z. Jaroszewicz<sup>1,4</sup>, M. Sarzyński<sup>5</sup>; <sup>1</sup>National Institute of Telecommunications (PL); <sup>2</sup>Warsaw University of Technology (PL); <sup>3</sup>Institute of Electronic Materials Technology (PL); <sup>4</sup>Institute of Applied Optics (PL); <sup>5</sup>Institute of High Pressure Physics (PL).*

A new method of manufacture of diffractive optical elements (DOEs) with variable diffraction efficiency is proposed. For this purpose the high energy beam sensitive (HEBS) glass is used as a halftone mask for generation of the binary phase gratings with variable phase step height in a resist layer. The main advantages as well as limitations of proposed technology are presented and discussed.

5009\_DO2012\_57

**Axicons with constant image intensity**

*Z. Jaroszewicz<sup>1,4</sup>, A. Kolodziejczyk<sup>2</sup>, M.S. Millan<sup>3</sup>, L.A. Romero<sup>3</sup>, M Sypek<sup>2</sup>, K. Petelczyc<sup>2</sup>; <sup>1</sup>National Institute of Telecommunications (PL); <sup>2</sup>Warsaw University of Technology (PL); <sup>3</sup>Technical University of Catalonia (ES); <sup>4</sup>Institute of Applied Optics (PL).*

We propose a new kind of axicons designed for the purposes of imaging with extended depth of focus (EDOF). They are distinguishable by the constant intensity of images produced within the whole range of the assumed foci lengths.

NOTES



LATE REGISTRATION FOR THE

8th EOS Topical Meeting on Diffractive Optics (DO 2012)

27 February - 1 March 2012 | Conference Centre of the Delft University of Technology, Delft, Netherlands  
[www.myeos.org/events/do2012](http://www.myeos.org/events/do2012)

Please return this form by fax to +49 (0)511 277 2699 or register online at [www.myeos.org/shop](http://www.myeos.org/shop)

REGISTRATION CATEGORY

The full-time-registration fee includes the participation in all four meeting days, one copy of the Topical Meeting digest CD-ROM, coffee breaks, and lunches on 27/28/29 February as well as the participation in the social programme on Wednesday, 29 February 2012. The participation in the following conference dinner is optional (additional costs tba).

Late registration (from 28 January 2012)	Universities, private persons, German and non-EU companies and non-university research institutes	EU companies and EU non-university research institutes (except Germany)*
Late registration for members	<input type="checkbox"/> 530.00 € (incl. 19% VAT)	<input type="checkbox"/> 445.38 €* (excl. VAT, VAT no. required)
Late registration for non-members	<input type="checkbox"/> 610.00 € (incl. 19% VAT)	<input type="checkbox"/> 512.61 €* (excl. VAT, VAT no. required)
Late registration for student members Copy of your current student ID required.	<input type="checkbox"/> 290.00 € (incl. 19% VAT)	<input type="checkbox"/> 243.70 €* (excl. VAT, VAT no. required)
Late registration for student non-members Copy of your current student ID required.	<input type="checkbox"/> 310.00 € (incl. 19% VAT)	<input type="checkbox"/> 260.50 €* (excl. VAT, VAT no. required)
Late registration for one-day No copy of digest CD-ROM included. You may directly order the digest below.  Please select the date of participation: <input type="checkbox"/> Monday, 27 Febr. <input type="checkbox"/> Tuesday, 28 Febr. <input type="checkbox"/> Wednesday, 29 Febr. <input type="checkbox"/> Thursday, 1 March	<input type="checkbox"/> 280.00 € (incl. 19% VAT)	<input type="checkbox"/> 235.29 €* (excl. VAT, VAT no. required)
Extra digest (CD-ROM)	<input type="checkbox"/> 40.00 € member rate (incl. 19% VAT) <input type="checkbox"/> 50.00 € non-member rate (incl. 19% VAT)	<input type="checkbox"/> 33.61 €* member rate (excl. 19% VAT, VAT no. required) <input type="checkbox"/> 42.02 €* non-member rate (excl. 19% VAT, VAT no. required)

I have read and accept the General Terms and Conditions of the EOS - Events & Services GmbH at: [http://www.myeos.org/aboutus/terms\\_and\\_conditions](http://www.myeos.org/aboutus/terms_and_conditions)

**\*IMPORTANT:** Registrations from companies and non-university research institutes registered in EU countries (except Germany) are exempted from VAT. Registrations without valid VAT no. will not be accepted. Please enter VAT no.:

VAT no.: \_\_\_\_\_



## LATE REGISTRATION FOR THE

## 8th EOS Topical Meeting on Diffractive Optics (DO 2012)

27 February - 1 March 2012 | Conference Centre  
of the Delft University of Technology, Delft, Netherlands  
[www.myeos.org/events/do2012](http://www.myeos.org/events/do2012)

Please return this form by fax to  
**+49 (0)511 277 2699**  
or register online at  
[www.myeos.org/shop](http://www.myeos.org/shop)

PLEASE FILL IN YOUR NAME AND ADDRESS

Page 2/2

Ms.  Mr. \_\_\_\_\_  
Title First name Name

Company

Street

City

Zip/Postal code

Country

Telephone

Fax

Email address

Homepage

## PAYMENT OPTIONS

## 1) CREDIT CARD PAYMENT

Payment shall be made in € by credit card. If bank transfer is preferred see details below.

Charge to my: Mastercard<sup>1</sup>  VISA<sup>1</sup>  Eurocard<sup>1</sup>  American Express<sup>2</sup>

Card No.:

□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

Expiry Date:

---

Verification No.:

---

<sup>[1]</sup>Mastercard, Visa, Eurocard: The final 3-digit number located on the back of your credit card.]

<sup>[2]</sup>American Express: The four small numbers printed on the front of your card, above the last few embossed numbers.]

Name of credit card holder:

---

Signature:

---

Date:

---

## 2) PAYMENT BY BANK TRANSFER

I wish to pay by bank transfer, please send me the invoice and banking details.

**Please note:** For payment via bank transfer we charge 15.00 € handling fee.

The full registration fee must arrive at the EOS bank account before the start of the meeting.

Please note that all bank fees must be paid by the attendee.

## CANCELLATION POLICY

Requests for cancellation must be made in writing to the EOS Office (address see below).

Refunds are subject to a processing fee of 50 €.

Cancellations received **by 6 February 2012:**

full refund minus processing fee (50 €).

Cancellations received **after 6 February 2012:**

no refund.



### **ANGE 2012**

#### **2nd Conference on Laser Ablation and Nanoparticle Generation in Liquids**

Hotel Caparena, Taormina (Sicily), Italy | 22 - 24 May 2012

[www.myeos.org/events/angel2012](http://www.myeos.org/events/angel2012) | [angel2012@myeos.org](mailto:angel2012@myeos.org)



### **TST 2012**

#### **3rd EOS Topical Meeting on Terahertz Science & Technology**

Prague, Czech Republic | 24 - 27 June 2012

[www.myeos.org/events/tst2012](http://www.myeos.org/events/tst2012) | [tst2012@myeos.org](mailto:tst2012@myeos.org)



### **AIT 2012**

#### **6th EOS Topical Meeting on Advanced Imaging Techniques**

Hyères - Southern Alps (Provence), France | 2 - 5 July 2012

[www.myeos.org/events/ait2012](http://www.myeos.org/events/ait2012) | [ait2012@myeos.org](mailto:ait2012@myeos.org)



### **EMVPO 2012**

#### **6th EOS Topical Meeting on Visual and Physiological Optics**

University College Dublin (UCD), Dublin, Ireland | 20 - 22 August 2012

[www.myeos.org/events/emvpo2012](http://www.myeos.org/events/emvpo2012) | [emvpo2012@myeos.org](mailto:emvpo2012@myeos.org)



### **EOSAM 2012**

#### **4th EOS Annual Meeting 2012**

Aberdeen Exhibition and Conference Centre, Aberdeen, Scotland | 25 - 28 September 2012

[www.myeos.org/events/eosam2012](http://www.myeos.org/events/eosam2012) | [aberdeen@myeos.org](mailto:aberdeen@myeos.org)

For more information about EOS organised and EOS co-sponsored events, please go to [www.myeos.org/events](http://www.myeos.org/events).

Should you wish to be kept informed about an EOS organised event, please send an email with your contact details to the above mentioned email address.

# Publish your research

## ....with JEOS:RP

Impact factor  
2010: 1.044

- open access journal
- professional peer-review
- listed with ISI Journal Citation Reports
- individual author support

### Editors

Joseph Braat, Delft University of Technology (NL)

Richard De La Rue, University of Glasgow (UK)

### Publication rates

350 € for EOS members\* | 400 € for non-members

\* Associate members are requested to upgrade to a full membership first (12.50 €/year).  
Special rates for attendees of EOS Events available! Check the event website for details.